

DOCKET NO.: 208671US0

TITLE OF THE INVENTION

BIS(2-ARYL-5-PYRIDYL) DERIVATIVES

5

BACKGROUND OF THE INVENTION

Field Of The Invention

10 The present invention relates to bis(2-aryl-5-pyridyl) derivatives or salts thereof, and to medicinal materials which comprise the bis(2-aryl-5-pyridyl) derivatives or salts thereof as active ingredients and are useful for the prevention or treatment of allergic immune diseases.

Discussion Of The Background

15 IgE, a class of immunoglobulin (Ig), is an allergen-specific molecule produced by IgE producing cells differentiated from B cells, when triggered by contact of immunocytes with an allergen in the body.

20 IgE is produced in a target organ exhibiting an allergic response, and binds to a receptor on the surfaces of mast cells, which are principal effector cells, in an allergic reaction or basophils (sensitized state). Allergic chemical mediators such as histamine, leucotrienes, prostaglandins, and PAF, and tissue destructive enzymes such as tryptase are released from the mast cells, which are stimulated as a result of intrusion of the allergen into the body after sensitization and its reaction with specific IgE, to provoke immediate responses of an allergic reaction such as increased vasopermeability, smooth muscle constriction and vasodilation. Cytokines, such as IL-4, which activate other immune system cells, are also secreted from the stimulated mast cells. As a result, eosinophils, basophils or

the like infiltrate tissues, and an allergic chemical mediator or tissue destructive protein such as MBP, which are secreted by these inflammatory cells, induce late responses of the allergic reaction and protract and worsen an allergic symptom.

As can be appreciated from what is stated above, abnormality in IgE production is highly relevant to various allergic immune diseases such as asthma, atopic dermatitis, allergic rhinitis, inflammatory bowel disease, contact dermatitis and allergic ophthalmopathy. It is known that inhibition of IgE production makes it possible to prevent and/or treat these diseases (Emerging Therapeutic Targets In Asthma And Allergy: Modulation Of IgE. Emerging Therapeutic Targets, 3, 229-240 (1990); Anti-IgE As Novel Therapy For The Treatment Of Asthma. Curr. Opin. Plum. Med., 5, 76-80 (1999); Treatment Of Allergic Asthma With Monoclonal Anti-IgE Antibody., N. Eng. J. Med., 341, 1966-1973 (1999); Anti-IgE Antibody Therapy For Asthma. N. Eng. J. Med., 341, 2006-2008 (1999)).


From the foregoing, IgE is believed to be a substance which is involved in the manifestation of an allergic disease from the very beginning. With the objective of developing antiallergic agents, some small molecules with IgE antibody production inhibiting activity have been found and have been reported to date (WO 98/04058, WO 98/07702, WO 98/16497, JP 10-324631A, WO 99/19291, WO 99/35140, WO 99/38829, WO 99/42446, JP 11-269192A, WO 00/05198, "Yakuri to Chiryō (Basic Pharmacology & Therapeutic)", 22(3), 1369 (1994), JP 1-106818A, JP 7-17506B, JP 8-92216A, JP 8-109177A, WO 96/11682, JP 59-167564A). These compounds, however, involve problems such as low solubility in water, and thus they are not fully satisfactory for the stated reason and other reasons.

#### SUMMARY OF THE INVENTION




Accordingly, one object of the present invention is to provide a compound having

excellent IgE antibody production inhibiting activity and also a medicinal composition comprising the compound as an active ingredient.

Briefly, this object and other objects of the present invention as hereinafter will become more readily apparent can be attained by a bis(2-aryl-5-pyridyl) derivative having  
 5 formula (1) or a salt thereof:



wherein A is a substituted or unsubstituted aromatic hydrocarbon group, and X is a group selected from one of the following formulas (2) to (4):



wherein in formula (2), R<sup>1</sup> is a hydrogen atom or a di(lower alkyl)amino(lower alkyl) group, and m is 1 or 2; in formula (3), Y<sup>1</sup> and Y<sup>2</sup> are each a nitrogen atom or a CH group, and n is 0 or an integer of 1 to 6; in formula (4), R<sup>2</sup> and R<sup>3</sup> each are a hydrogen atom or a lower alkyl group, Z is a single bond, a substituted methylene group, a substituted imino group, an oxygen atom or a cycloalkylene group, and p and q are each 0 or an integer of 1 to 6.

Another aspect of the invention is a medicinal composition comprising the bis(2-aryl-5-pyridyl) derivative or the salt thereof and a pharmacologically acceptable carrier.

Still another aspect of the invention is a method of treating a subject for an allergic

immune disease by administering the bis(2-aryl-5-pyridyl) derivative or the salt thereof to a subject.

#### DESCRIPTION OF THE PREFERRED EMBODIMENTS

5 As a result of investigation, it has now been found that compounds represented by formula (1) have excellent IgE antibody production inhibiting activity and also good solubility in water and hence, are useful as medicines for the prevention or treatment of allergic immune diseases.

10 Illustrative of embodiments of lower alkyl in "lower alkyl groups", "lower alkoxy groups", "halogeno(lower alkoxy) groups", "lower alkoxy(lower alkoxy) groups", "hydroxy(lower alkoxy) groups", "(lower alkoxy)carbonyl groups", "lower alkanoyl groups", "lower alkanoyloxy groups", "(lower alkyl)- and/or (lower alkoxy)-substituted carbamoyl groups", "(lower alkyl)thio groups", "(lower alkyl)amino groups", "(lower alkyl)sulfonylamino groups" as used herein include linear, branched or cyclic alkyl groups  
15 having 1 to 6 carbon atoms. Suitable examples include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, cyclopropyl, and cyclohexyl. Suitable examples of "halogen atoms" are fluorine, chlorine, bromine, and iodine.

*Ammon A<sup>1</sup>*  
20 ~~In formula (1), the aromatic hydrocarbon group represented by A is preferably a group which has 6 to 14 carbon atoms, with a phenyl or naphthyl group being more preferred and a phenyl group being particularly preferred. These groups may contain 1 to 3 substituents. Suitable examples of such substituents include lower alkyl groups, lower alkoxy groups, halogeno(lower alkyl) groups, lower alkoxy(lower alkyl) groups, hydroxy(lower alkyl) groups, carboxyl group, (lower alkoxy)carbonyl groups, unsubstituted or (lower alkyl)- and/or (lower alkoxy)-substituted carbamoyl groups, lower alkanoyl groups,~~

a<sup>1</sup>  
cont.  
formyl group, lower alkanoyloxy groups, halogen atoms, hydroxy group, cyano, (lower alkyl)thio groups, amino group, mono- or di-(lower alkyl)amino groups, (lower alkyl)sulfonylamino groups, pyrrolidinyl groups, and alkylenedioxy groups.

Preferred specific examples of these substituents include methyl, methoxy, ethoxy, n-propoxy, isopropoxy, trifluoroethoxy, methoxyethoxy, hydroxyethoxy, hydroxy, cyano, methylthio, dimethylamino, pyrrolidinyl, carboxyl, ethoxycarbonyl, t-butoxycarbonyl, butyryloxy, N-methyl-N-methoxycarbamoyl, acetyl, and methylenedioxy.

Among the groups represented by X, the group of formula (2) contains the di(lower alkyl)amino(lower alkyl) group represented by R<sup>1</sup>. As the di(lower alkyl)amino group in the di(lower alkyl)amino(lower alkyl) group, a dimethylamino group or the like is particularly preferred.

In formula (3), n preferably ranges from 2 to 4.

In formula (4), the substituted methylene group represented by Z contains one or two substituents, examples of which include lower alkyl, hydroxy, pyrrolidinyl and benzyloxy groups. The substituted imino group represented by Z contains one or two substituents, examples of which include lower alkyl groups. Illustrative of the cycloalkylene represented by Z are cyclopentylene, cyclohexylene, and cycloheptylene.


Further, p and q preferably range from 0 to 4.

Preferred specific examples of bis(2-aryl-5-pyridyl) derivative (1) of the present invention include N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine dimethanesulfonate, N,N'-bis[2-(4-hydroxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine dimethanesulfonate, 1,4-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]hexahydro-1,4-diazepine dimethanesulfonate, N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-diethylethylenediamine dimethanesulfonate, N,N'-

bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine  
 dimethanesulfonate, N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N',2,2-tetramethyl-  
 1,3-propanediamine dimethanesulfonate, 2-hydroxy-N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-  
 pyridyl]-N,N'-dimethyl-1,3-propanediamine, N,N-bis[N-[2-(3,4,5-trimethoxyphenyl)-5-  
 5 pyridyl]-N-methyl-2-aminoethyl]methylamine trihydrochloride, N,N-bis[N-[2-(3,4,5-  
 trimethoxyphenyl)-5-pyridyl]-N-methyl-3-aminopropyl]methylamine, bis[N-[2-(3,4,5-  
 trimethoxyphenyl)-5-pyridyl]-2-aminoethyl]ether dimethanesulfonate, 2-  
 dimethylaminomethyl-1,4-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]piperazine  
 trihydrochloride, N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,6-  
 hexanediamine dimethanesulfonate, N,N-bis[N-[2-[3,5-dimethoxy-4-(2,2,2-  
 trifluoroethoxy)phenyl]-5-pyridyl]-N-methyl-3-aminopropyl]methylamine trihydrochloride.

Compound (1) of the present invention can be produced, for example, by the reaction  
 steps of the below-described processes, although no particular limitation is imposed on the  
 process of synthesis employed.

( Process 1)



wherein A and X have the same meanings as defined above, W and W' represent halogen or

-OSO<sub>2</sub>(C<sub>r</sub>F<sub>2r+1</sub>) in which r is 0 or an integer of 1 to 4, and M represents dihydroxyboron, di(lower alkoxy)boron, di(lower alkyl)boron, dihalo(lower alkyl)silicon, halogenated zinc, tri(lower alkyl)tin, halogenated magnesium or the like.

Described specifically, compound (1) of the present invention can be prepared by adding compound (6) and a catalyst to a solution or suspension of compound (5), conducting the reaction optionally in the presence of a ligand and a base to prepare compound (7) and then treating compound (7) with compound (8).

#### 1) Synthesis of the compound (7) (cross-coupling reaction)

Compound (7) can be obtained by adding compound (6), the catalyst and, if necessary, the ligand and the base to the solution or suspension of compound (5) and then conducting the reaction at room temperature to 200°C for 0.5 to 10 hours (Metal-catalyzed Cross-coupling Reactions; Diederich, F., Stang, P.J., Eds.; Wiley-VHC: Weinheim (1998). Stanforth, S.P. Tetrahedron, 54, 263-303 (1998)).

Illustrative of a solvent usable in the above reaction include benzene, toluene, xylene, diethyl ether, tetrahydrofuran, dimethoxyethane, dioxane, acetonitrile, dimethylformamide, N-methylpiperidone, methanol, ethanol, and water. Examples of the catalyst include tetrakis(triphenylphosphine)palladium(0), tris(bisbenzylideneacetone)dipalladium(0), palladium(II) acetate, palladium(II) chloride, dichlorobis(triphenylphosphine)palladium(II), dichloro[1,2-bis(diphenylphosphino)ethane]palladium(II), dichloro[1,4-bis(diphenylphosphino)butane]palladium(II), dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium(II), tetrakis(triphenylphosphine)nickel(0) and bis(acetylacetonato)nickel (II).

Suitable examples of the ligand of the metal complexes include tri(t-butyl)phosphine, triphenylphosphine, tri(o-tolyl)phosphine, tri(2-furyl)phosphine, 2,2'-bis(diphenyl-

phosphino)-1,1'-binaphthyl, 1,2-bis(diphenylphosphino)-ethane, 1,3-bis(diphenylphosphino)propane, 1,4-bis(diphenylphosphino)butane, and 1,1'-bis(diphenylphosphino)ferrocene. Suitable examples of a base include sodium acetate, potassium carbonate, sodium carbonate, sodium hydrogencarbonate, potassium phosphate, sodium hydroxide, potassium hydroxide, barium hydroxide, sodium methoxide, sodium ethoxide, cesium fluoride, tetrabutylammonium fluoride, and triethylamine.

The reaction can be conducted preferably by following the procedure disclosed by Stille et al (Stille, J.K., Angew. Chem. Int. Ed. Engl., 25, 508-524 (1986).), the procedure disclosed by Suzuki et al (Miyaura, N.; Suzuki, A. Chem. Rev., 95, 2457-2483 (1995)) or the procedure disclosed by Mitchell et al, (Mitchell, M.B.; Wallbank, P.J., Tetrahedron Lett., 32, 2273-2276 (1991)). That is, compound (6) in which M is a tri(lower alkyl)tin can be used under conditions of tetrakis(triphenylphosphine)palladium(0)/toluene/100 to 150°C/10 to 30 hours. Also compound (6) in which M is a dihydroxyboron can be used under conditions of tetrakis(triphenylphosphine)palladium(0)/sodium carbonate (or potassium carbonate)/water-ethanol(or methanol)-toluene/60 to 100°C/0.5 to 3 hours. Still further, compound (6) in which M is a dihydroxyboron can be used under conditions of dichloro[1,2-bis(diphenylphosphino)ethane]palladium(II)[or dichloro[1,4-bis(diphenylphosphino)butane]palladium(II)]/sodium carbonate(or potassium carbonate)/water-ethanol(or methanol)-toluene/80 to 120°C/1 to 24 hours.

## 2) Synthesis of the compound (1) (condensation)

Compound (1) may also be obtained by adding compound (8), a catalyst, a base and, if necessary, a ligand to a solution of compound (7) and then conducting the reaction at room temperature to 200°C for 0.5 to 100 hours, preferably at 80 to 120°C for 5 to 15 hours (Yang, B.H., Buchwald, S.L., J. Organomet. Chem., 576, 125-146 (1999). Hartwig, J.F., Angew.



Chem. Int. Ed. Engl., 37, 2046-2067 (1998). Nishiyama, M., Yamamoto, T., Koike, Y.,  
Tetrahedron Lett., 39, 617-620 (1998)).


Suitable solvents for the reaction include toluene, xylene, dimethoxyethane, tetrahydrofuran, and dioxane. Suitable catalyst embodiments include

5 tetrakis(triphenylphosphine)palladium(0), tris(bisbenzylideneacetone)dipalladium(0), palladium(II) acetate, and dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium(II), with tetrakis(triphenylphosphine)palladium(0) and tris(bisbenzylideneacetone)dipalladium(0) being particularly preferred. Suitable embodiments of the base include sodium t-butoxide, potassium t-butoxide, potassium phosphate, potassium carbonate, cesium carbonate, lithium

10 tetramethyldisilazide, triethylamine, and 1,8-diazabicyclo[5,4,0]-7-undecene(DBU), with sodium t-butoxide, potassium t-butoxide and potassium carbonate being particularly preferred. Suitable examples of the ligand include tri(t-butyl)phosphine, triphenylphosphine, tri(o-tolyl)phosphine, tri(2-furyl)phosphine, 2,2'-bis(diphenyl-phosphino)-1,1'-binaphthyl, 1,2-bis(diphenylphosphino)ethane, 1,3-bis(diphenylphosphino)propane, 1,4-

15 bis(diphenylphosphino)butane, and 1,1'-bis(diphenyl-phosphino)ferrocene.

(Production Process 2)



wherein A, X, W, W' and M have the same meanings as defined above.

More specifically, compound (1) can be prepared by subjecting compound (9) and compound (8) to a condensation reaction thereby preparing compound (10), removing the benzyl group of compound (10) to yield compound (11), modifying the hydroxyl groups of compound (11) to prepare compound (12), and then treating compound (12) with compound (6).

#### 1) Synthesis of the compound (10) (condensation)

The compound (10) may be prepared by treating 2-benzyloxy-5-halopyridine (9) with compound (8) under similar conditions as in the reaction of compound (7) and compound (8) in Process 1 above.

#### 2) Synthesis of the compound (11) (removal of the benzyl group)

Compound (11) may be prepared by adding an acid to a solution of compound (10) and then conducting the reaction at -20 to 100°C for 0.5 to 100 hours, preferably at 30 to 60°C for 1 to 6 hours.

Illustrative of suitable solvents include chloroform, methylene chloride, dichloroethane, and ethyl acetate. Suitable examples of the acid include hydrogen chloride and trifluoroacetic acid. It is also possible to use the acid as a solvent.

3) Synthesis of the compound (12) (conversion of the hydroxyl group into a better leaving group)

Compound (12) can be prepared by adding a base and  $\text{ClSO}_2(\text{C}_5\text{F}_{2s+1})$  or  $[\text{SO}_2(\text{C}_5\text{F}_{2s+1})]_2\text{O}$  (being  $[\text{SO}_2(\text{CF}_3)]_2\text{O}$  preferred) (s is 0 or an integer of 1 to 4) to a solution or suspension of compound (11) and stirring the resulting mixture at  $-78$  to  $100^\circ\text{C}$  for 5 minutes to 10 hours, preferably at  $-78$  to  $0^\circ\text{C}$  for 10 minutes to 2 hours.

Illustrative of suitable solvents include methylene chloride, chloroform, dichloroethane, benzene, toluene, xylene, diethyl ether, tetrahydrofuran, dimethoxyethane, dioxane, and acetonitrile. Suitable examples of the base include sodium acetate, potassium carbonate, sodium hydrogencarbonate, potassium phosphate, triethylamine, N,N-diisopropylethylamine, and pyridine, with triethylamine and N,N-diisopropylethylamine being preferred.

In an alternative process, compound (12) can be prepared by treating compound (11) with a halogenating reagent at room temperature to  $200^\circ$  for 0.5 minutes to 200 hours, preferably at  $50$  to  $120^\circ\text{C}$  for 5 to 12 hours. A solvent can also be used as needed.


Suitable halogenating reagents include phosphorus pentachloride, phosphorus oxychloride, phosphorus tribromide, and phosphorus oxybromide, with phosphorus oxychloride-phosphorus pentachloride being particularly preferred. Suitable examples of the solvent include chloroform, methylene chloride and dichloroethane.

4) Synthesis of the compound (1) (cross-coupling reaction)

Compound (1) can be prepared by treating compound (12) with compound (6) under similar conditions as in the cross-coupling of compound (5) with compound (6) in Process 1

above.

(Production Process 3)



wherein A, W, M,  $R^2$  and  $R^3$  have the same meanings as defined above, and  $R^4$  represents a lower alkyl group.

Among compounds (1), compounds (1') in which X is a group of formula (4), p and q are 3, and Z is a substituted imino group can be prepared by the reaction steps of the above-described Process 3.

1) Synthesis of compound (14) (cross-coupling reaction)

Compound (14) can be prepared by treating 2-halo-5-nitropyridine (13) with compound (6) under similar conditions as described in the cross-coupling reaction between compound (5) and compound (6) in Process 1 above.

2) Synthesis of compound (15) (reduction of the nitro group)

Compound (15) can be prepared by reducing compound (14) in a manner known *per*

se in the art. As a specific reducing reaction, a catalyst and a hydrogen source are added to a solution of compound (14), followed by conducting the reaction at 0 to 100°C for 5 minutes to 100 hours, preferably at 20 to 60°C for 0.5 to 5 hours (Process A); or an acid and a metal or metal salt are added to a solution or suspension of compound (14), followed by conducting the reaction at room temperature to 100°C for 5 minutes to 100 hours, preferably at 50 to 100°C for 0.5 to 3 hours (Process B).

For Process A, suitable solvents include tetrahydrofuran, dimethoxyethane, dioxane, ethyl acetate, methanol, ethanol. Suitable examples of catalysts include palladium on charcoal, platinum black, platinum, and Raney nickel; and illustrative sources of hydrogen include hydrogen, ammonium formate, and hydrazine.

As representative of specific conditions for Process B, suitable solvents include ethanol and water. Suitable examples of acids include hydrochloric acid and acetic acid; and illustrative examples of metals or metal salts include zinc powder, tin powder, iron powder, and stannous chloride.

### 3) Synthesis of compound (16) (acylation)

Compound (16) can be prepared by adding a base and acryloyl chloride to a solution of compound (15) and treating them at -78 to 50°C for 5 minutes to 100 hours, preferably at -30 to 30°C for 10 minutes to 1 hour.

Suitable solvents include methylene chloride, chloroform, dichloroethane, benzene, toluene, xylene, diethyl ether, tetrahydrofuran, dimethoxyethane, dioxane, and acetonitrile, with methylene chloride and tetrahydrofuran being particularly preferred. Suitable bases include sodium acetate, potassium carbonate, sodium hydrogencarbonate, potassium phosphate, triethylamine, and N,N-diisopropylethylamine, with triethylamine and N,N-diisopropylethylamine being particularly preferred.

4) Synthesis of compound (18) (Michael addition)

Compound (18) can be prepared by treating compound (16) with amine (17) at 0 to 200°C for 0.5 to 100 hours, preferably at 20 to 100°C for 1 to 15 hours.

Here, a solvent may be used if necessary. Suitable solvents include methylene chloride, chloroform, dichloroethane, benzene, toluene, xylene, diethyl ether, tetrahydrofuran, dimethoxyethane, dioxane, acetonitrile, and dimethylformamide.

5) Synthesis of compound (19) (reduction)

Compound (19) can be prepared by adding a reducing agent to a solution of compound (18) and treating the prepared material at 0 to 100°C for 5 minutes to 100 hours, preferably at 20 to 80°C for 30 minutes to 3 hours.

Suitable solvents include toluene, diethyl ether, tetrahydrofuran, and dimethoxyethane. Suitable reducing agents include lithium aluminum hydride, diisobutylaluminum hydride, and borane.

6) Synthesis of compound (1') (alkylating reaction)

Compound (1') can be prepared by adding a reducing agent and an acid to a solution of compound (19) and a carbonyl compound and conducting the reaction at -20 to 50°C for 0.5 to 100 hours, preferably at 0 to 30°C for 1 to 15 hours.

Suitable carbonyl compounds include formaldehyde, lower alkylaldehydes, and di(lower alkyl) ketone. Suitable solvents include methylene chloride, chloroform, tetrahydrofuran, dimethoxyethane, dioxane, acetonitrile, methanol, and ethanol.

Suitable reducing agents include lithium aluminum hydride, sodium borohydride, sodium cyanoborohydride, and borane. Suitable acids include hydrochloric acid, acetic acid, and trifluoroacetic acid.

(Process 4):



solvent, in the presence of a coupling agent, in the presence of a base added as needed, at 0 to 100°C for 0.5 to 30 hours (Process A); or intermediate (15) and a reactive derivative of dicarboxylic acid (20) are treated in a solvent, in the presence of a base added as needed, at 0 to 100°C for 0.5 to 30 hours (Process B). In each of Process A and Process B, the reaction is preferably conducted at 0 to 50°C for 0.5 to 5 hours.

Illustrative of suitable solvents employed in these reactions include dimethylformamide, tetrahydrofuran, dioxane, acetonitrile, methylene chloride, and dichloroethane. Illustrative of bases include organic bases such as pyridine, triethylamine, and N,N-diisopropylethylamine; and inorganic bases such as sodium carbonate and sodium hydrogencarbonate.

Further, examples of the coupling agent used in the process A include 1,3-dicyclohexylcarbodiimide, 1-cyclohexyl-3-morpholinoethylcarbodiimide, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide, 1,1'-carbonyldiimidazole, diethyl phosphorocyanidate, diphenylphosphoryl azide, bis(2-oxo-3-oxazolidinyl)phosphonic chloride, and 2-chloro-1-methylpyridinium iodide. Suitable examples of the reactive derivative of the dicarboxylic acid (20) in Process B include acid halides such as acid chlorides, acid azides, mixed anhydrides with pivalic acid or the like, and activated esters such as cyanomethyl esters and p-nitrophenyl esters.

## 2) Synthesis of compound (1'') (reduction)

Compound (1'') can be prepared by treating compound (21) under similar conditions as in the reduction of compound (18) in Process 3.

## 3) Synthesis of compound (1''') (alkylation)

Compound (1''') can be prepared by treating compound (1'') under similar conditions as in the alkylation of compound (19) in Process 3.



05693580-052901  
The intermediate and objective compounds prepared in the above reactions, respectively, can be isolated and purified by subjecting them to a purification procedure commonly employed in organic synthetic chemistry, for example, filtration, extraction, washing, drying, concentration, recrystallization, one or more of various types of chromatography, and the like. Further, the intermediate can be provided for use in the next reaction without specifically purifying the intermediate.

In addition, the intermediates and objective compounds may also be prepared in the form of solvates with reaction solvents, recrystallization solvents or the like, especially as hydrates. Further, compound (1) of the present invention may include various isomers depending on the kinds and combination of the substituents. It is to be noted that the present invention encompasses all of such isomers.

Compound (1) obtained as described above can be converted into an acid addition salt or a basic salt by a method known *per se* in the art. No particular limitation is imposed on such salts insofar as they are pharmacologically acceptable salts. When compound (1) is a basic compound, examples of pharmacologically acceptable salt include mineral acid salts such as the hydrochloride, sulfate and nitrate; and organic acid salts such as the methanesulfonate, acetate, oxalate and citrate. When compound (1) is an acidic compound, on the other hand, examples of its pharmacologically acceptable salt include alkali metal salts such as the sodium and potassium salts; alkaline earth metal salts such as the calcium and magnesium salts; and organic base salts such as the pyridine, picoline and triethylamine salts.

The bis (2-aryl-5-pyridyl) derivative (1) of the present invention has excellent IgE antibody production inhibiting activity as will be demonstrated in tests to be described subsequently herein and also, IL-4 production inhibiting activity and IL-5 production inhibiting activity. The compound is also useful as a medicinal agent for the prevention or

treatment of various allergic diseases, for example, asthma, atopic dermatitis, allergic rhinitis, inflammatory bowel disease, contact skin dermatitis and allergic ophthalmopathy, and also as an IgE antibody production inhibitor.

5 The medicinal agent of the present invention comprises, as an active ingredient, the bis(2-aryl-5-pyridyl) derivative or the salt thereof. By adding pharmacologically acceptable, inorganic or organic carriers, the bis(2-aryl-5-pyridyl) derivative or the salt thereof can be formulated into medicinal compositions, for example, various oral preparations or parenteral preparations such as solid, semi-solid or liquid preparations by methods known *per se* in the art.

Illustrative of preparations for oral administration include tablets, pills, granules, soft or hard capsules, triturates, subtilized granules, powders, emulsions, syrups, pellets, and elixirs. On the other hand, illustrative of preparations for parenteral administration include injections, drips, infusions, ointments, lotions, tonics, sprays, suspensions, medicinal oils, emulsions, suppositories, and instillations.

To formulate such preparations, methods known *per se* in the art can be followed. The active ingredient of the present invention can be used in combination with pharmacologically acceptable surfactants, excipients, coloring matters, flavoring agents, preservatives, stabilizers, buffering agents, suspending agents, isotonicities and the like as needed.

20 The dosage of the medicinal agent of the present invention varies *inter alia* depending on the medicine, the target disease to be treated or prevented, the method of administration, the treatment period, and the age, sex and weight of the patient. Nonetheless, the medicinal agent is preferably administered at a daily dosage ranging from 0.01 to 1,000 mg/kg · weight in terms of the compound represented by formula (1). This dosage can be


administered at once or in several portions, for example, 2 to 6 portions in a day.

Having now generally described this invention, a further understanding can be obtained by reference to certain specific examples which are provided herein for purposes of illustration only and are not intended to be limiting unless otherwise specified.

5

#### Reference Example 1

N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine



Under argon, tri-*t*-butylphosphine (5.0 mL, 20 mmol), sodium *t*-butoxide (10.78 g, 112 mmol), 2-benzyloxy-5-bromopyridine [*J. Am. Chem. Soc.*, **71**, 70-73 (1949).] (29.20 g, 110 mmol) and a solution of N,N'-dimethylethylenediamine (4.40 g, 50.0 mmol) in *o*-xylene (30 mL) were added to a solution of bis(dibenzylideneacetone)palladium (2.86 g, 5.00 mmol) in *o*-xylene (80 mL). After stirring at 120°C for 3 hours, water was added, and the resulting mixture was extracted with chloroform. The organic layer was washed with water, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to afford N,N'-bis(2-benzyloxy-5-pyridyl)-N,N'-dimethylethylenediamine as a pale yellow crystalline powder (melting point: 95.0-97.0°C)(18.59 g, yield: 81%).

A solution of N,N'-bis(2-benzyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (18.50 g, 400 mmol) in trifluoroacetic acid (105 mL) was stirred at 50°C for 4 hours. Diethyl ether (630 mL) was added to the ice-cold reaction mixture, and the mixture was stirred for 1 hour.


Precipitated crystals were collected by filtration and then washed with diethyl ether to give N,N'-bis(2-hydroxy-5-pyridyl)-N,N'-dimethylethylenediamine ditrifluoroacetate as a yellow

crystalline powder (melting point: 180.0-182.0°C)(17.24 g, yield 86%).

N,N'-Bis(2-hydroxy-5-pyridyl)-N,N'-dimethylethylenediamine ditrifluoroacetate (18.00 g, 35.8 mmol) was suspended in methylene chloride (350 mL) and N,N-diisopropylethylamine (48 mL, 280 mmol) was added. The resulting mixture was cooled to -10°C, and trifluoromethanesulfonic anhydride (21 mL, 120 mmol) was added. After stirring for 30 minutes, N,N-diisopropylethylamine (12 mL, 70 mmol) and trifluoromethanesulfonic anhydride (21 mL, 120 mmol) were added. After being stirred further at -10°C for 30 minutes, N,N-diisopropylethylamine (5.0 mL, 29 mmol) and trifluoromethanesulfonic anhydride (5.0 mL, 28 mmol) were added additionally, and the mixture was stirred for 1 hour in an ice bath. A saturated aqueous sodium carbonate (180 mL) and sodium carbonate (18 g, 0.17 mmol) were added, and the resulting mixture was extracted with methylene chloride. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to obtain crude crystals. The crude crystals were recrystallized from chloroform-hexane to afford the title compound as pale brown needles (melting point: 146.0-148.0°C )(13.44 g, yield: 70%).

#### Reference Example 2

##### 5-Chloro-2-(3,4,5-trimethoxyphenyl)pyridine




Under argon, 3,4,5-trimethoxyphenylboronic acid (5.05 g, 24.0 mmol), dicyclo[1,2-bis(diphenylphosphino)ethane]palladium (574.0 mg, 1.00 mmol) and a 2.0 M aqueous

contd.  
Q2

sodium carbonate (20 mL, 40 mmol) were added to a solution of 2,5-dichloropyridine (2.95 g, 20.0 mmol) in ethanol-toluene (1:5, 54.0 mL). After stirring at 100°C for 3 hours, water was added, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to afford the title compound as a colorless crystalline powder (melting point: 61.0-63.0°C)(4.13 g, yield: 73%).

### Reference Example 3

#### 4-Ethoxy-3,5-dimethoxyphenylboronic acid




To a solution of 4-iodo-2,6-dimethoxyphenol (JP61-207432A) (100.0 mg, 0.360 mmol) in dimethylformamide (1.0 mL) were added ethyl iodide (84.0 mg, 0.540 mmol) and a 50% dispersion of sodium hydride in mineral oil (19.0 mg, 0.40 mmol). After stirring at room temperature for 1 hour, the reaction mixture was poured into 0.10 M hydrochloric acid, and the resulting mixture was extracted with diethyl ether. The organic layer was washed with saturated aqueous sodium hydrogencarbonate and brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to afford 4-ethoxy-1-iodo-3,5-dimethoxybenzene as a colorless amorphous powder (102.0 mg, yield: 93%).

Under argon, a 1.60 M solution of n-butyllithium in hexane (2.5 mL, 4.1 mmol) was added to anhydrous tetrahydrofuran (7.0 mL) stirred in a dry ice-acetone bath, followed by dropwise addition of a solution of 4-ethoxy-1-iodo-3,5-dimethoxybenzene (570.0 mg, 1.85 mmol) in anhydrous tetrahydrofuran (1.5 mL). After the mixture was stirred for 20

minutes in the dry ice-acetone bath, triisopropyl borate (0.50 mL, 2.2 mmol) was added, and the mixture was additionally stirred for 20 minutes. The reaction mixture was stirred at room temperature for 1 hour, and concentrated under reduced pressure. To the residue, a 0.1 M aqueous sodium hydroxide (4.4 mL) solution was added. The aqueous solution was washed with chloroform, acidified by the addition of concentrated hydrochloric acid, and extracted with chloroform. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. Chloroform-hexane was added to the residue and the resulting precipitate was collected by filtration to afford the title compound as a colorless amorphous powder (162.0 mg, yield: 39%).

#### Reference Example 4

##### 3,5-Dimethoxy-4-(2,2,2-trifluoroethoxy)phenylboronic acid



To a solution of methyl syringate (2.10 g, 9.90 mmol) in dimethylformamide (21.0 mL) were added potassium carbonate (13.60 g, 9.90 mmol) and 2,2,2-trifluoroethyl trifluoromethanesulfonate (purity: 25%) (*J. Org. Chem.*, 30, 4322-4324 (1965)) (10.90 g, 12 mmol).

After stirring at room temperature for 1 hour and at 50°C for 3 hours, water was added and the resulting mixture was extracted with diethyl ether. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to prepare an oil (2.69 g) containing methyl 3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)benzoate.

To a solution of the oil (2.69 g) obtained in methanol (14 mL) was added a 5.0 M aqueous sodium hydroxide (14 mL, 70 mmol), and the resulting mixture was stirred at 80°C for 30 minutes. The reaction mixture was concentrated under reduced pressure to remove the

methanol, acidified by the addition of concentrated hydrochloric acid, and extracted with chloroform. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to yield an oil (2.56 g) containing 3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)benzoic acid.

5 To a solution of the oil (2.45 g) prepared by the above-described procedure in t-butyl alcohol (37 mL) were added triethylamine (0.93 g, 9.2 mmol) and diphenylphosphoryl azide (2.53 g, 9.20 mmol). After stirring at 100°C for 2 hours, the reaction mixture was concentrated under reduced pressure. A solution of the residue in ethyl acetate was washed with saturated aqueous sodium hydrogencarbonate and brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to provide N-t-butoxycarbonyl-3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)aniline as a colorless oil (1.10 g, yield: 36% based on methyl syringate).


10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995  
1000  
1005  
1010  
1015  
1020  
1025  
1030  
1035  
1040  
1045  
1050  
1055  
1060  
1065  
1070  
1075  
1080  
1085  
1090  
1095  
1100  
1105  
1110  
1115  
1120  
1125  
1130  
1135  
1140  
1145  
1150  
1155  
1160  
1165  
1170  
1175  
1180  
1185  
1190  
1195  
1200  
1205  
1210  
1215  
1220  
1225  
1230  
1235  
1240  
1245  
1250  
1255  
1260  
1265  
1270  
1275  
1280  
1285  
1290  
1295  
1300  
1305  
1310  
1315  
1320  
1325  
1330  
1335  
1340  
1345  
1350  
1355  
1360  
1365  
1370  
1375  
1380  
1385  
1390  
1395  
1400  
1405  
1410  
1415  
1420  
1425  
1430  
1435  
1440  
1445  
1450  
1455  
1460  
1465  
1470  
1475  
1480  
1485  
1490  
1495  
1500  
1505  
1510  
1515  
1520  
1525  
1530  
1535  
1540  
1545  
1550  
1555  
1560  
1565  
1570  
1575  
1580  
1585  
1590  
1595  
1600  
1605  
1610  
1615  
1620  
1625  
1630  
1635  
1640  
1645  
1650  
1655  
1660  
1665  
1670  
1675  
1680  
1685  
1690  
1695  
1700  
1705  
1710  
1715  
1720  
1725  
1730  
1735  
1740  
1745  
1750  
1755  
1760  
1765  
1770  
1775  
1780  
1785  
1790  
1795  
1800  
1805  
1810  
1815  
1820  
1825  
1830  
1835  
1840  
1845  
1850  
1855  
1860  
1865  
1870  
1875  
1880  
1885  
1890  
1895  
1900  
1905  
1910  
1915  
1920  
1925  
1930  
1935  
1940  
1945  
1950  
1955  
1960  
1965  
1970  
1975  
1980  
1985  
1990  
1995  
2000  
2005  
2010  
2015  
2020  
2025  
2030  
2035  
2040  
2045  
2050  
2055  
2060  
2065  
2070  
2075  
2080  
2085  
2090  
2095  
2100  
2105  
2110  
2115  
2120  
2125  
2130  
2135  
2140  
2145  
2150  
2155  
2160  
2165  
2170  
2175  
2180  
2185  
2190  
2195  
2200  
2205  
2210  
2215  
2220  
2225  
2230  
2235  
2240  
2245  
2250  
2255  
2260  
2265  
2270  
2275  
2280  
2285  
2290  
2295  
2300  
2305  
2310  
2315  
2320  
2325  
2330  
2335  
2340  
2345  
2350  
2355  
2360  
2365  
2370  
2375  
2380  
2385  
2390  
2395  
2400  
2405  
2410  
2415  
2420  
2425  
2430  
2435  
2440  
2445  
2450  
2455  
2460  
2465  
2470  
2475  
2480  
2485  
2490  
2495  
2500  
2505  
2510  
2515  
2520  
2525  
2530  
2535  
2540  
2545  
2550  
2555  
2560  
2565  
2570  
2575  
2580  
2585  
2590  
2595  
2600  
2605  
2610  
2615  
2620  
2625  
2630  
2635  
2640  
2645  
2650  
2655  
2660  
2665  
2670  
2675  
2680  
2685  
2690  
2695  
2700  
2705  
2710  
2715  
2720  
2725  
2730  
2735  
2740  
2745  
2750  
2755  
2760  
2765  
2770  
2775  
2780  
2785  
2790  
2795  
2800  
2805  
2810  
2815  
2820  
2825  
2830  
2835  
2840  
2845  
2850  
2855  
2860  
2865  
2870  
2875  
2880  
2885  
2890  
2895  
2900  
2905  
2910  
2915  
2920  
2925  
2930  
2935  
2940  
2945  
2950  
2955  
2960  
2965  
2970  
2975  
2980  
2985  
2990  
2995  
3000  
3005  
3010  
3015  
3020  
3025  
3030  
3035  
3040  
3045  
3050  
3055  
3060  
3065  
3070  
3075  
3080  
3085  
3090  
3095  
3100  
3105  
3110  
3115  
3120  
3125  
3130  
3135  
3140  
3145  
3150  
3155  
3160  
3165  
3170  
3175  
3180  
3185  
3190  
3195  
3200  
3205  
3210  
3215  
3220  
3225  
3230  
3235  
3240  
3245  
3250  
3255  
3260  
3265  
3270  
3275  
3280  
3285  
3290  
3295  
3300  
3305  
3310  
3315  
3320  
3325  
3330  
3335  
3340  
3345  
3350  
3355  
3360  
3365  
3370  
3375  
3380  
3385  
3390  
3395  
3400  
3405  
3410  
3415  
3420  
3425  
3430  
3435  
3440  
3445  
3450  
3455  
3460  
3465  
3470  
3475  
3480  
3485  
3490  
3495  
3500  
3505  
3510  
3515  
3520  
3525  
3530  
3535  
3540  
3545  
3550  
3555  
3560  
3565  
3570  
3575  
3580  
3585  
3590  
3595  
3600  
3605  
3610  
3615  
3620  
3625  
3630  
3635  
3640  
3645  
3650  
3655  
3660  
3665  
3670  
3675  
3680  
3685  
3690  
3695  
3700  
3705  
3710  
3715  
3720  
3725  
3730  
3735  
3740  
3745  
3750  
3755  
3760  
3765  
3770  
3775  
3780  
3785  
3790  
3795  
3800  
3805  
3810  
3815  
3820  
3825  
3830  
3835  
3840  
3845  
3850  
3855  
3860  
3865  
3870  
3875  
3880  
3885  
3890  
3895  
3900  
3905  
3910  
3915  
3920  
3925  
3930  
3935  
3940  
3945  
3950  
3955  
3960  
3965  
3970  
3975  
3980  
3985  
3990  
3995  
4000  
4005  
4010  
4015  
4020  
4025  
4030  
4035  
4040  
4045  
4050  
4055  
4060  
4065  
4070  
4075  
4080  
4085  
4090  
4095  
4100  
4105  
4110  
4115  
4120  
4125  
4130  
4135  
4140  
4145  
4150  
4155  
4160  
4165  
4170  
4175  
4180  
4185  
4190  
4195  
4200  
4205  
4210  
4215  
4220  
4225  
4230  
4235  
4240  
4245  
4250  
4255  
4260  
4265  
4270  
4275  
4280  
4285  
4290  
4295  
4300  
4305  
4310  
4315  
4320  
4325  
4330  
4335  
4340  
4345  
4350  
4355  
4360  
4365  
4370  
4375  
4380  
4385  
4390  
4395  
4400  
4405  
4410  
4415  
4420  
4425  
4430  
4435  
4440  
4445  
4450  
4455  
4460  
4465  
4470  
4475  
4480  
4485  
4490  
4495  
4500  
4505  
4510  
4515  
4520  
4525  
4530  
4535  
4540  
4545  
4550  
4555  
4560  
4565  
4570  
4575  
4580  
4585  
4590  
4595  
4600  
4605  
4610  
4615  
4620  
4625  
4630  
4635  
4640  
4645  
4650  
4655  
4660  
4665  
4670  
4675  
4680  
4685  
4690  
4695  
4700  
4705  
4710  
4715  
4720  
4725  
4730  
4735  
4740  
4745  
4750  
4755  
4760  
4765  
4770  
4775  
4780  
4785  
4790  
4795  
4800  
4805  
4810  
4815  
4820  
4825  
4830  
4835  
4840  
4845  
4850  
4855  
4860  
4865  
4870  
4875  
4880  
4885  
4890  
4895  
4900  
4905  
4910  
4915  
4920  
4925  
4930  
4935  
4940  
4945  
4950  
4955  
4960  
4965  
4970  
4975  
4980  
4985  
4990  
4995  
5000  
5005  
5010  
5015  
5020  
5025  
5030  
5035  
5040  
5045  
5050  
5055  
5060  
5065  
5070  
5075  
5080  
5085  
5090  
5095  
5100  
5105  
5110  
5115  
5120  
5125  
5130  
5135  
5140  
5145  
5150  
5155  
5160  
5165  
5170  
5175  
5180  
5185  
5190  
5195  
5200  
5205  
5210  
5215  
5220  
5225  
5230  
5235  
5240  
5245  
5250  
5255  
5260  
5265  
5270  
5275  
5280  
5285  
5290  
5295  
5300  
5305  
5310  
5315  
5320  
5325  
5330  
5335  
5340  
5345  
5350  
5355  
5360  
5365  
5370  
5375  
5380  
5385  
5390  
5395  
5400  
5405  
5410  
5415  
5420  
5425  
5430  
5435  
5440  
5445  
5450  
5455  
5460  
5465  
5470  
5475  
5480  
5485  
5490  
5495  
5500  
5505  
5510  
5515  
5520  
5525  
5530  
5535  
5540  
5545  
5550  
5555  
5560  
5565  
5570  
5575  
5580  
5585  
5590  
5595  
5600  
5605  
5610  
5615  
5620  
5625  
5630  
5635  
5640  
5645  
5650  
5655  
5660  
5665  
5670  
5675  
5680  
5685  
5690  
5695  
5700  
5705  
5710  
5715  
5720  
5725  
5730  
5735  
5740  
5745  
5750  
5755  
5760  
5765  
5770  
5775  
5780  
5785  
5790  
5795  
5800  
5805  
5810  
5815  
5820  
5825  
5830  
5835  
5840  
5845  
5850  
5855  
5860  
5865  
5870  
5875  
5880  
5885  
5890  
5895  
5900  
5905  
5910  
5915  
5920  
5925  
5930  
5935  
5940  
5945  
5950  
5955  
5960  
5965  
5970  
5975  
5980  
5985  
5990  
5995  
6000  
6005  
6010  
6015  
6020  
6025  
6030  
6035  
6040  
6045  
6050  
6055  
6060  
6065  
6070  
6075  
6080  
6085  
6090  
6095  
6100  
6105  
6110  
6115  
6120  
6125  
6130  
6135  
6140  
6145  
6150  
6155  
6160  
6165  
6170  
6175  
6180  
6185  
6190  
6195  
6200  
6205  
6210  
6215  
6220  
6225  
6230  
6235  
6240  
6245  
6250  
6255  
6260  
6265  
6270  
6275  
6280  
6285  
6290  
6295  
6300  
6305  
6310  
6315  
6320  
6325  
6330  
6335  
6340  
6345  
6350  
6355  
6360  
6365  
6370  
6375  
6380  
6385  
6390  
6395  
6400  
6405  
6410  
6415  
6420  
6425  
6430  
6435  
6440  
6445  
6450  
6455  
6460  
6465  
6470  
6475  
6480  
6485  
6490  
6495  
6500  
6505  
6510  
6515  
6520  
6525  
6530  
6535  
6540  
6545  
6550  
6555  
6560  
6565  
6570  
6575  
6580  
6585  
6590  
6595  
6600  
6605  
6610  
6615  
6620  
6625  
6630  
6635  
6640  
6645  
6650  
6655  
6660  
6665  
6670  
6675  
6680  
6685  
6690  
6695  
6700  
6705  
6710  
6715  
6720  
6725  
6730  
6735  
6740  
6745  
6750  
6755  
6760  
6765  
6770  
6775  
6780  
6785  
6790  
6795  
6800  
6805  
6810  
6815  
6820  
6825  
6830  
6835  
6840  
6845  
6850  
6855  
6860  
6865  
6870  
6875  
6880  
6885  
6890  
6895  
6900  
6905  
6910  
6915  
6920  
6925  
6930  
6935  
6940  
6945  
6950  
6955  
6960  
6965  
6970  
6975  
6980  
6985  
6990  
6995  
7000  
7005  
7010  
7015  
7020  
7025  
7030  
7035  
7040  
7045  
7050  
7055  
7060  
7065  
7070  
7075  
7080  
7085  
7090  
7095  
7100  
7105  
7110  
7115  
7120  
7125  
7130  
7135  
7140  
7145  
7150  
7155  
7160  
7165  
7170  
7175  
7180  
7185  
7190  
7195  
7200  
7205  
7210  
7215  
7220  
7225  
7230  
7235  
7240  
7245  
7250  
7255  
7260  
7265  
7270  
7275  
7280  
7285  
7290  
7295  
7300  
7305  
7310  
7315  
7320  
7325  
7330  
7335  
7340  
7345  
7350  
7355  
7360  
7365  
7370  
7375  
7380  
7385  
7390  
7395  
7400  
7405  
7410  
7415  
7420  
7425  
7430  
7435  
7440  
7445  
7450  
7455  
7460  
7465  
7470  
7475  
7480  
7485  
7490  
7495  
7500  
7505  
7510  
7515  
7520  
7525  
7530  
7535  
7540  
7545  
7550  
7555  
7560  
7565  
7570  
7575  
7580  
7585  
7590  
7595  
7600  
7605  
7610  
7615  
7620  
7625  
7630  
7635  
7640  
7645  
7650  
7655  
7660  
7665  
7670  
7675  
7680  
7685  
7690  
7695  
7700  
7705  
7710  
7715  
7720  
7725  
7730  
7735  
7740  
7745  
7750  
7755  
7760  
7765  
7770  
7775  
7780  
7785  
7790  
7795  
7800  
7805  
7810  
7815  
7820  
7825  
7830  
7835  
7840  
7845  
7850  
7855  
7860  
7865  
7870  
7875  
7880  
7885  
7890  
7895  
7900  
7905  
7910  
7915  
7920  
7925  
7930  
7935  
7940  
7945  
7950  
7955  
7960  
7965  
7970  
7975  
7980  
7985  
7990  
7995  
8000  
8005  
8010  
8015  
8020  
8025  
8030  
8035  
8040  
8045  
8050  
8055  
8060  
8065  
8070  
8075  
8080  
8085  
8090  
8095  
8100  
8105  
8110  
8115  
8120  
8125  
8130  
8135  
8140  
8145  
8150  
8155  
8160  
8165  
8170  
8175  
8180  
8185  
8190  
8195  
8200  
8205  
8210  
8215  
8220  
8225  
8230  
8235  
8240  
8245  
8250  
8255  
8260  
8265  
8270  
8275  
8280  
8285  
8290  
8295  
8300  
8305  
8310  
8315  
8320  
8325  
8330  
8335  
8340  
8345  
8350  
8355  
8360  
8365  
8370  
8375  
8380  
8385  
8390  
8395  
8400  
8405  
8410  
8415  
8420  
8425  
8430  
8435  
8440  
8445  
8450  
8455  
8460  
8465  
8470  
8475  
8480  
8485  
8490  
8495  
8500  
8505  
8510  
8515  
8520  
8525  
8530  
8535  
8540  
8545  
8550  
8555  
8560  
8565  
8570  
8575  
8580  
8585  
8590  
8595  
8600  
8605  
8610  
8615  
8620  
8625  
8630  
8635  
8640  
8645  
8650  
8655  
8660  
8665  
8670  
8675  
8680  
8685  
8690  
8695  
8700  
8705  
8710  
8715  
8720  
8725  
8730  
8735  
8740  
8745  
8750  
8755  
8760  
8765  
8770  
8775  
8780  
8785  
8790  
8795  
8800  
8805  
8810  
8815  
8820  
8825  
8830  
8835  
8840  
8845  
8850  
8855  
8860  
8865  
8870  
8875  
8880  
8885  
8890  
8895  
8900  
8905  
8910  
8915  
8920  
8925  
8930  
8935  
8940  
8945  
8950  
8955  
8960  
8965  
8970  
8975  
8980  
8985  
8990  
8995  
9000  
9005  
9010  
9015  
9020  
9025  
9030  
9035  
9040  
9045  
9050  
9055  
9060  
9065  
9070  
9075  
9080  
9085  
9090  
9095  
9100  
9105  
9110  
9115  
9120  
9125  
9130  
9135  
9140  
9145  
9150  
9155  
9160  
9165  
9170  
9175  
9180  
9185  
9190  
9195  
9200  
9205  
9210  
9215  
9220  
9225  
9230  
9235  
9240  
9245  
9250  
9255  
9260  
9265  
9270  
9275  
9280  
9285  
9290  
9295  
9300  
9305  
9310  
9315  
9320  
9325  
9330  
9335  
9340  
9345  
9350  
9355  
9360  
9365  
9370  
9375  
9380  
9385  
9390  
9395  
9400  
9405  
9410  
9415  
9420  
9425  
9430  
9435  
9440  
9445  
9450  
9455  
9460  
9465  
9470  
9475  
9480  
9485  
9490  
9495  
9500  
9505  
9510  
9515  
9520  
9525  
9530  
9535  
9540  
9545  
9550  
9555  
9560  
9565  
9570  
9575  
9580  
9585  
9590  
9595  
9600  
9605  
9610  
9615  
9620  
9625  
9630  
9635  
9640  
9645  
9650  
9655  
9660  
9665  
9670  
9675  
9680  
9685  
9690  
9695  
9700  
9705  
9710  
9715  
9720  
9725  
9730  
9735  
9740  
9745  
9750  
9755  
9760  
9765  
9770  
9775  
9780  
9785  
9790  
9795  
9800  
9805  
9810  
9815  
9820  
9825  
9830  
9835  
9840  
9845  
9850  
9855  
9860  
9865  
9870  
9875  
9880  
9885  
9890  
9895  
9900  
9905  
9910  
9915  
9920  
9925  
9930  
9935  
9940  
9945  
9950  
9955  
9960  
9965  
9970  
9975  
9980  
9985  
9990  
9995  
10000  
10005  
10010  
10015  
10020  
10025  
10030  
10035  
10040  
10045  
10050  
10055  
10060  
10065  
10070  
10075  
10080  
10085  
10090  
10095  
10100  
10105  
10110  
10115  
10120  
10125  
10130  
10135  
10140  
10145  
10150  
10155  
10160

organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield 1-iodo-3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)benzene as a colorless oil [1.00 g, yield: 88% based on N-t-butoxy-carbonyl-3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)aniline].

5 Applying the synthetic procedure of 4-ethoxy-3,5-dimethoxyphenylboronic acid from 4-ethoxy-1-iodo-3,5-dimethoxybenzene in Reference Example 3, the title compound was prepared as a colorless amorphous powder (369.0 mg, yield: 48%) from 1-iodo-3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)-benzene (1.00 g, 2.76 mmol).

#### Reference Example 5

##### 4-(2-Methoxyethoxy)-3,5-dimethoxyphenylboronic acid



To a solution of 4-iodo-2,6-dimethoxyphenol (JP61-207432A) (280.0 mg, 1.00 mmol) in dimethylformamide (3.0 mL) were added potassium carbonate (184.0 mg, 1.10 mmol) and ethyl bromoacetate (184.0 mg, 1.10 mmol). After the resulting mixture was stirred at room temperature for 1 hour and at 50°C for 1 hour, water was added, and resulting mixture was extracted with diethyl ether. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to afford an oil (351.0 mg) containing 4-ethoxy-carbonylmethoxy-1-iodo-3,5-dimethoxybenzene.

20 Lithium aluminum hydride (11.0 mg, 0.29 mmol) was added to an ice-cold solution of the oil (155.0 mg, approx. 0.42 mmol) prepared by the above-described procedure in anhydrous diethyl ether (2.0 mL), and the mixture was stirred for 10 minutes. After the resulting mixture was stirred at room temperature for 1 hour, 3.0 M hydrochloric acid was




added, and the aqueous layer was extracted with diethyl ether. The organic layer was washed with saturated aqueous sodium hydrogencarbonate and brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to afford 4-(2-hydroxyethoxy)-1-iodo-3,5-dimethoxybenzene as a colorless oil (83.0 mg, yield: 59% based on 4-iodo-2,6-dimethoxyphenol).

Methyl iodide (0.010 mL, 0.16 mmol) and a 50% dispersion of sodium hydride in mineral oil (12.0 mg, 0.24 mmol) were added to a solution of 4-(2-hydroxyethoxy)-1-iodo-3,5-dimethoxybenzene (39.0 mg, 0.120 mmol) in anhydrous tetrahydrofuran (1.0 mL), and the mixture was stirred at room temperature for 1 hour. Water was added to the reaction mixture, and the mixture was extracted with diethyl ether. The organic layer was washed with 3.0 M hydrochloric acid, saturated aqueous sodium hydrogencarbonate and brine, dried over anhydrous sodium sulfate, and then concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield 1-iodo-3,5-dimethoxy-4-(2-methoxyethoxy)benzene as a colorless oil (40.0 mg, yield: 98%).

Applying the synthetic procedure of 4-ethoxy-3,5-dimethoxyphenylboronic acid from 4-ethoxy-1-iodo-3,5-dimethoxybenzene in Reference Example 3, the title compound was obtained as a colorless amorphous powder (276.0 mg, yield: 54%) from 1-iodo-3,5-dimethoxy-4-(2-methoxyethoxy)benzene (673.0 mg, 2.00 mmol).

#### Reference Example 6

4-[2-(*t*-butyldimethylsiloxy)ethoxy]-3,5-dimethoxyphenylboronic acid




To a solution of 4-(2-hydroxyethoxy)-1-iodo-3,5-dimethoxybenzene (2.76 g, 8.52 mmol) synthesized by the process shown in Reference Example 5 in dimethylformamide (25 mL) were added imidazole (1.16 g, 17.0 mmol) and t-butylchlorodimethylsilane (1.61 g, 10.7 mmol), and the mixture was stirred at 50°C for 3 hours. Water was added to the reaction mixture, and the mixture was extracted with ethyl acetate-hexane (1:1). The organic layer was washed with water and brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield 4-[2-(t-butyltrimethylsiloxy)-ethoxy]-1-iodo-3,5-dimethoxybenzene as a colorless oil (3.56 g, yield: 95%).

Applying the synthetic procedure of 4-ethoxy-3,5-dimethoxyphenylboronic acid from 4-ethoxy-1-iodo-3,5-dimethoxybenzene in Reference Example 3, the title compound was obtained as a colorless amorphous powder (1.15 g; quantitative) from 4-[2-(t-butyltrimethylsiloxy)ethoxy]-1-iodo-3,5-dimethoxybenzene (1.42 g, 3.24 mmol).

#### Reference Example 7

##### 3,5-Dimethoxy-4-(N-methoxy-N-methylcarbamoyl)phenylboronic acid




To a solution of 4-iodo-2,6-dimethoxybenzoic acid (308.0 mg, 1.00 mmol) in methylene chloride (5.0 mL) were added dimethylformamide (0.020 mL, 0.26 mmol) and oxalyl chloride (0.13 mL, 1.5 mmol), and the resulting mixture was stirred for 30 minutes. The reaction mixture was poured into an ice-cold solution of N,O-dimethylhydroxylamine hydrochloride (488.0 mg, 5.00 mmol) and N,N-diisopropylethylamine (1.74 mL, 10.0 mmol) in methylene chloride (5.0 mL). The ice bath was removed, and the mixture was stirred at

room temperature for 30 minutes. 0.1 M Hydrochloric acid was added to the reaction mixture, and the resulting mixture was extracted with chloroform. The organic layer was washed with 0.1 M aqueous sodium hydroxide and brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. Hexane was added to the residue and the resulting precipitate was collected by filtration to yield crude 4-iodo-N,2,6-trimethoxy-N-methylbenzamide as a pale yellow crystalline powder (305.0 mg, yield: approx. 87%).

Applying the synthetic procedure of 4-ethoxy-3,5-dimethoxyphenylboronic acid from 4-ethoxy-1-iodo-3,5-dimethoxybenzene in Reference Example 3, the title compound was obtained as a pale yellow amorphous powder (91.0 mg, yield: 36% based on 4-iodo-2,6-dimethoxybenzoic acid) from the crude 4-iodo-N,2,6-trimethoxy-N-methylbenzamide (287.0 mg, approx. 0.82 mmol).

#### Reference Example 8

(4-Acetyl-3,5-dimethoxyphenyl)trimethyltin



Aluminum chloride (87.05 g, 653.0 mmol) was gradually added to a solution of 3,5-dimethoxyaniline (20.0 g, 131.0 mmol) and acetic anhydride (30.9 mL, 328 mmol) in methylene chloride (650 mL) stirred in a sodium chloride-ice bath. The resulting mixture was stirred in the sodium chloride-ice bath for 20 minutes and at room temperature for 1 hour. The reaction mixture was poured into ice water, and the mixture was extracted with chloroform. The organic layer was washed with saturated aqueous sodium hydrogencarbonate and brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to

prepare crude crystals. The crude crystals were recrystallized from diethyl ether-hexane to yield 4-acetamido-2,6-dimethoxyacetophenone as a pale yellow crystalline powder (melting point 184.0-187.0°C )(9.42 g, yield: 26%).

To a suspension of 4-acetamido-2,6-dimethoxyacetophenone (9.42 g, 39.7 mmol) in ethanol (400 mL) was added a 5.0 M aqueous sodium hydroxide solution (80 mL, 400 mL) and the mixture was stirred at 95°C for 3 hours. The reaction mixture was concentrated under reduced pressure, water was added to the residue, and the resulting mixture was extracted with chloroform. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was suspended in diethyl ether-hexane (1:4) and collected by filtration to yield 4-amino-2,6-dimethoxyacetophenone as a pale yellow crystalline powder (melting point: 155.0-158.0°C)(7.58 g, yield: 98%).


To a suspension of 4-amino-2,6-dimethoxyacetophenone (7.58 g, 38.8 mmol) in water (155 mL) stirred in a sodium chloride-ice bath was added concentrated hydrochloric acid (6.5 mL, 78 mmol) to provide a homogeneous solution. To the solution stirred in the sodium chloride-ice bath was added a solution of sodium nitrite (2.81 g, 40.7 mmol) in water (8.0 mL) gradually and the resulting mixture was stirred in an ice bath for 15 minutes. A solution of potassium iodide (7.08 g, 42.7 mmol) in water (8.0 mL) and ethyl acetate (48 mL) were added, and the mixture was stirred at room temperature for 1 hour and at 50°C for 15 minutes. After cooling, the aqueous layer was extracted with ethyl acetate. The organic layers were combined, washed with saturated aqueous sodium hydrogencarbonate, a 10% aqueous sodium thiosulfate and brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield 4-iodo-2,6-dimethoxyacetophenone, obtained as a colorless crystalline powder

(melting point: 124.0-129.0°C )(9.50 g, yield: 80%).

Under nitrogen, to a solution of 4-iodo-2,6-dimethoxyacetophenone (287.0 mg, 1.86 mmol) in anhydrous 1,4-dioxane (12 mL) were added hexamethylditin (609.0 mg, 1.86 mmol) and tetrakis(triphenylphosphine)palladium (215.0 mg, 0.190 mmol), and the mixture was stirred at 110°C for 4 hours. After cooling, water was added and the mixture was extracted with benzene. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield the title compound as a colorless oil (450.0 mg, yield: 71%).

#### Reference Example 9


N,N'-Bis(2-chloro-5-pyridyl)-N,N'-dimethylethylenediamine



To phosphorus oxychloride (16.4 mL) were added N,N'-bis(2-hydroxy-5-pyridyl)-N,N'-dimethylethylenediamine ditrifluoroacetate (824.0 mg, 1.64 mmol) synthesized by the method described in Reference Example 1 and phosphorus pentachloride (171.0 mg, 0.820 mmol). The resulting mixture was stirred at 100°C for 8 hours. After cooling, the reaction mixture was poured into ice water. The mixture was rendered basic by adding sodium hydroxide and then extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford crude crystals of the title compound as a pale brown crystalline powder (melting point: 173.0-179.0°C)(439.0 mg, yield: approx. 86%).

#### Reference Example 10

2-Benzoyloxy-N,N'-dimethyl-1,3-propanediamine



To a solution of 2-hydroxy-1,3-propanediamine (0.90 g, 10 mmol) in tetrahydrofuran (40 mL) was added triethylamine (3.94 g, 39 mmol). To the ice-cold solution was added a solution of ethyl chloroformate (3.58 g, 33 mmol) in tetrahydrofuran (10 mL) dropwise. The resulting mixture was stirred at room temperature for 15 hours, and filtered to remove insoluble materials. The filtrate was dried over anhydrous sodium sulfate and concentrated under reduced pressure to yield an oil (1.95 g) containing N,N'-bis(ethoxycarbonyl)-2-ethoxycarbonyloxy-1,3-propanediamine.

Lithium aluminum hydride (780.0 mg, 20.7 mmol) was added to a solution of the oil (1.95 g, approx. 6.5 mmol) prepared by the above-described procedure in tetrahydrofuran (40 mL). An ice bath was removed, and the mixture was stirred at 80°C for 17 hours. After cooling, anhydrous sodium sulfate (25.0 g) was added to the reaction mixture. Insoluble materials were removed by filtration through Celite, and the filtrate was concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to give 2-hydroxy-N,N'-dimethyl-1,3-propanediamine as a pale yellow oil (277.0 mg, yield: 23% based on 2-hydroxy-1,3-propanediamine).

To a solution of 2-hydroxy-N,N'-dimethyl-1,3-propanediamine (165.0 mg, 1.39 mmol) in acetonitrile (5.0 mL) were added triethylamine (564.0 mg, 5.58 mmol) and di-*t*-butyl dicarbonate (912.0 mg, 4.18 mmol). After stirring at room temperature for 15 hours, 0.10 M hydrochloric acid was added and the mixture was extracted with chloroform. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel


to provide N,N'-bis(t-butoxycarbonyl)-2-hydroxy-N,N'-dimethyl-1,3-propanediamine as a pale yellow oil (163.0 mg, yield: 36%).

To an ice-cold solution of N,N'-bis(t-butoxy-carbonyl)-2-hydroxy-N,N'-dimethyl-1,3-propanediamine (87.0 mg, 0.27 mmol) in tetrahydrofuran (4.0 mL) were added benzyl bromide (230.0 mg, 1.35 mmol) and a 50% dispersion of sodium hydride in mineral oil (35.0 mg, 0.81 mmol). The reaction mixture was stirred at room temperature for 15 hours and concentrated under reduced pressure. Water was added to the residue, and the mixture was extracted with ethyl acetate. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield 2-benzyloxy-N,N'-bis(t-butoxycarbonyl)-N,N'-dimethyl-1,3-propanediamine, obtained as a colorless oil (54.0 mg, yield: 49%).

To a solution of 2-benzyloxy-N,N'-bis(t-butoxycarbonyl)-N,N'-dimethyl-1,3-propanediamine (73.0 mg, 0.17 mmol) in methylene chloride (2.0 mL) was added a 4.0 M solution of hydrogen chloride in ethyl acetate (1.0 mL, 4.0 mmol), and the mixture was stirred at room temperature for 4 hours. The reaction mixture was concentrated under reduced pressure, ethanol (5.0 mL) was added to the residue, and the resulting mixture was concentrated under reduced pressure. An approx. 10% solution of ammonia in methanol (2.0 mL, approx. 12 mmol) was added to the residue, insoluble materials were removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield the title compound as a colorless oil (35.0 mg, yield: 98%).

Reference Example 11

5-Amino-2-(3,4,5-trimethoxyphenyl)pyridine




Under nitrogen, tetrakis(triphenylphosphine)palladium (504.6mg, 0.437 mmol) and a 2.0 M aqueous sodium carbonate (25 mL, 50 mmol) were added to a solution of 2-bromo-5-nitropyridine (2.87 g, 14.1 mmol) and 3,4,5-trimethoxyphenylboronic acid (3.00 g, 14.2 mmol) in ethanol-toluene (1:5, 90 mL), and the resulting mixture was stirred at 70°C for 12 hours. Water was added to the reaction mixture, and the aqueous layer was extracted with ethyl acetate. The organic layers were combined, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel and crude crystals were obtained. The crude crystals were recrystallized from chloroform-hexane to provide 5-nitro-2-(3,4,5-trimethoxyphenyl)pyridine as yellow needles (melting point: 134.0-136.5°C )(3.31 g, yield: 81%).

To a solution of 5-nitro-2-(3,4,5-trimethoxyphenyl)-pyridine (1.50 g, 5.17 mmol) in methanol-ethyl acetate (1:4, 50 mL) was added 10% palladium on charcoal (158.6 mg), and the resulting mixture was stirred at 45°C for 2 hours and 30 minutes under hydrogen. Insoluble materials were removed by filtration through Celite, and the filtrate was concentrated under reduced pressure. The residue was recrystallized from ethyl acetate-hexane to yield the title compound as colorless needles (melting point: 153.5-155.5°C )(1.25 g, yield: 93%).

Reference Example 12

N,N'-Dimethyl-3-(1-pyrrolidinyl)-1,5-pentanediamine





Following the process described in the publication [*J. Am. Chem. Soc.*, **110**, 6127-6179 (1988)], N,N'-dibenzyl-N,N'-dimethyl-3-(1-pyrrolidinyl)glutaramide was prepared as a colorless oil (2.81 g, yield: 90% based on 6-chloro-2-pyrone) from 6-chloro-2-pyrone (1.00 g, 7.66 mmol) and N-methylbenzylamine (3.71 g, 30.6 mmol) via crude N,N'-dibenzyl-N,N'-dimethylglutaconamide (2.67 g).


Under nitrogen, lithium aluminum hydride (0.44 g, 12 mmol) was added to a solution of N,N'-dibenzyl-N,N'-dimethyl-3-(1-pyrrolidinyl)glutaramide (1.16 g, 2.85 mmol) in anhydrous tetrahydrofuran (20 mL) stirred in an ice bath, and the mixture was stirred at 65°C for 2 hours. Methanol (3.0 mL, 74 mmol) was added to the ice-cold reaction mixture, and the ice bath was removed. Water (2.0 mL), diethyl ether (150 mL) and anhydrous magnesium sulfate (15 g) were added, and the mixture was stirred at room temperature for 1 hour. Insoluble materials were removed by filtration through Celite, and the filtrate was concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield N,N'-dibenzyl-N,N'-dimethyl-3-(1-pyrrolidinyl)-1,5-pentanediamine as a colorless oil (917.3 mg, yield: 85%).

To a solution of N,N'-dibenzyl-N,N'-dimethyl-3-(1-pyrrolidinyl)-1,5-pentanediamine (249.1 mg, 0.657 mmol) in methanol (3.0 mL) were added concentrated hydrochloric acid (0.33 mL, 4.0 mmol) and 10% palladium on charcoal (50.0 mg), and the mixture was stirred at 50°C for 14 hours under hydrogen. Insoluble materials were removed by filtration through a membrane filter, and the filtrate was concentrated under reduced pressure. The residue was adsorbed on an alumina column, and eluted with methanol-chloroform (1:4). The eluate was concentrated under reduced pressure to afford the title compound as a

colorless oil (101.9 mg, yield: 78%).

### Example 1

N,N'-Bis[2-(2,4-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine  
dimethanesulfonate




To a solution of N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (108.0 mg, 0.200 mmol) synthesized in Reference Example 1 in ethanol-toluene (1:6, 3.5 mL) were added 2,4-dimethoxyphenylboronic acid (108.0 mg, 0.200 mmol), tetrakis-(triphenylphosphine)palladium (46.0 mg, 0.040 mmol) and a 2.0 M aqueous sodium carbonate solution (0.20 mL, 0.40 mmol). After stirring at 80°C for 2 hours, brine was added, and the mixture was extracted with chloroform. The organic layer was washed with brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified column chromatography on silica gel to provide N,N'-bis[2-(2,4-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine as a colorless amorphous powder (109.0 mg; quantitative).

To a solution of N,N'-bis[2-(2,4-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (108.0 mg, 0.200 mmol) in methanol (2.0 mL) was added a 1.0 M solution of methanesulfonic acid in methanol (0.44 mL, 0.44 mmol), and the reaction mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to yield the title compound as a yellow crystalline powder (melting point: 195.0-197.0°C)(61.0 mg, yield: 43%).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 120°C) (ammonium salt NH<sup>+</sup> protons were not observed) δ: 2.38(s, 6H), 3.07(s, 6H), 3.75(s, 4H), 3.85(s, 12H), 6.70(dd, J=2.4, 8.5Hz, 2H), 6.74(d, J=2.4Hz, 2H), 7.51(d, J=8.5Hz, 2H), 7.69(dd, J=2.9, 9.2Hz, 2H), 7.82(d, J=9.2Hz, 2H), 8.01(d, J=2.9Hz, 2H).

## Example 2

N,N'-Bis[2-(2,3-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine dimethanesulfonate




Following the procedure of Example 1, N,N'-bis[2-(2,3-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine was prepared as a colorless oil (33.0 mg, yield: 31%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (114.0 mg, 0.212 mmol) synthesized as in Reference Example 1 and 2,3-dimethoxyphenylboronic acid (128.0 mg; 0.850 mmol).

To a solution of N,N'-bis[2-(2,3-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (33.0 mg, 0.0641 mmol) in chloroform (3.0 mL) was added a 1.0 M solution of methanesulfonic acid in methanol (0.13 mL, 0.13 mmol), and the reaction mixture was concentrated under reduced pressure. Methanol-diethyl ether was added to the residue and the resulting precipitate was collected by filtration to yield the title compound as a yellow crystalline powder (melting point: 191.5-194.5°C)(39.5 mg, yield: 87%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)(ammonium salt NH<sup>+</sup> protons were not observed) δ: 2.85(s, 6H), 3.39(s, 6H), 3.79(s, 6H), 3.83(s, 4H), 3.92(s, 6H), 7.03-7.40(m, 6H), 7.61(br d, J=9.2Hz, 2H), 7.97(d, J=9.2Hz, 2H), 8.95(br s, 2H).

### 5 Example 3

N,N'-Bis[2-(2,6-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine  
dimethanesulfonate



Following the procedure of Example 1, crude crystals were obtained from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (150.0 mg, 0.279 mmol) synthesized in Reference Example 1 and 2,6-dimethoxyphenylboronic acid (150.7 mg, 1.00 mmol). The crude crystals were recrystallized from chloroform-diethyl ether-hexane to yield N,N'-bis[2-(2,6-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine as a colorless crystalline powder (77.3 mg, yield: 54%).

To a solution of N,N'-bis[2-(2,6-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (72.3 mg, 0.141 mmol) in methanol-chloroform (9:1, 10 mL) was added a 1.0 M aqueous methanesulfonic acid solution (0.35 mL, 0.35 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from ethanol-diethyl ether to provide the title compound as a pale yellow crystalline powder (melting point: 243.0-245.0°C)(94.1 mg, yield: 95%).


<sup>1</sup>H-NMR (CDCl<sub>3</sub>)(ammonium salt NH<sup>+</sup> protons were not observed) δ: 2.79(s, 6H),

3.28(s, 6H), 3.79(s, 4H), 3.83(s, 12H),  
 6.65(d, J=8.5Hz, 4H), 7.39(t, J=8.5Hz, 2H),  
 7.52(br dd, J=3.1, 9.2Hz, 2H),  
 7.69(br d, J=9.2Hz, 2H), 8.91(br d, J=3.1Hz, 2H).

5

#### Example 4

*N,N'*-Bis[2-(3,5-dimethoxyphenyl)-5-pyridyl]-*N,N'*-dimethylethylenediamine



Following the procedure of Example 1, an oil was obtained from *N,N'*-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-*N,N'*-dimethylethylenediamine (108.0 mg, 0.200 mmol) synthesized in Reference Example 1 and 2,5-dimethoxyphenylboronic acid (91.0 mg, 0.500 mmol). Chloroform-diethyl ether-hexane was added to the oil and the resulting precipitate was collected by filtration to yield the title compound as a pale yellow crystalline powder (melting point: 139.0-141.0°C) (41.2 mg, yield: 40%).


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 3.02(s, 6H), 3.66(s, 4H), 3.87(s, 12H),  
 6.46(br s, 2H), 6.99(br d, J=9.2Hz, 2H),  
 7.11(br s, 4H), 7.57(br d, J=9.2Hz, 2H),  
 8.22(br s, 2H).

15

20

#### Example 5

*N,N'*-Bis[2-(2,3,5-trimethoxyphenyl)-5-pyridyl]-*N,N'*-dimethylethylenediamine




Following the procedure of Example 1, crude crystals were obtained from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (150.0 mg, 0.280 mmol) synthesized in Reference Example 1 and 2,3,5-trimethoxyphenylboronic acid (131.0 mg, 0.620 mmol). The crude crystals were recrystallized from chloroform-hexane to yield the title compound as pale brown needles (melting point: 171.0-172.0°C) (37.0 mg, yield: 56%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 3.05(s, 6H), 3.60(s, 6H), 3.67(s, 4H), 3.85(s, 6H), 3.88(s, 6H), 6.50(d, J=2.9Hz, 2H), 6.95(d, J=2.9Hz, 2H), 7.01(dd, J=2.9, 8.8Hz, 2H), 7.85(d, J=8.8Hz, 2H), 8.25(d, J=2.9Hz, 2H).

#### Example 6

N,N'-Bis[2-(2,4,6-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine dimethanesulfonate



Following the procedure of Example 1, N,N'-bis[2-(2,4,6-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine was obtained as a colorless oil (20.1 mg, yield: 13%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (150.3 mg, 0.279 mmol) synthesized in Reference Example 1 and 2,4,6-


trimethoxyphenylboronic acid (124.5 mg, 0.692 mmol).

To a solution of N,N'-bis[2-(2,4,6-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (20.1 mg, 0.0350 mmol) in methanol-chloroform (1:1, 5 mL) was added a 1.0 M aqueous methanesulfonic acid solution (0.090 mL, 0.090 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to yield the title compound as pale yellow fine flakes (melting point: 264.0-265.0°C) (15.6 mg, yield: 59%).

<sup>1</sup>H-NMR [CD<sub>3</sub>OD-CDCl<sub>3</sub>(1:20)](ammonium salt NH<sup>+</sup> protons were not observed) δ: 2.77(s, 6H), 3.19(s, 6H), 3.80(s, 4H), 3.82(s, 12H), 3.88(s, 6H), 6.21(s, 4H), 7.59(dd, J=3.1, 9.3Hz, 2H), 7.69(d, J=9.3Hz, 2H), 8.50(dd, J=3.1Hz, 2H).

#### Example 7

N,N'-Bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine dimethanesulfonate



Under argon, tri-*t*-butylphosphine (101.0 mg, 0.500 mmol) was added to a solution of bis(dibenzylideneacetone)palladium (71.0 mg, 0.120 mmol) in *o*-xylene (3.0 mL). To the resulting solution were added 5-chloro-2-(3,4,5-trimethoxyphenyl)pyridine (335.0 mg, 1.20 mmol) synthesized in Reference Example 2, N,N'-dimethylethylenediamine (44.0 mg, 0.500

mmol), and sodium t-butoxide (106.0 mg, 1.10 mmol). After stirring at 120°C for 12 hours, water was added to the reaction mixture and the aqueous layer was extracted with chloroform. The organic layers were combined, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine as a pale yellow amorphous powder (167.0 mg, yield: 58%).


To a solution of N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (140.0 mg, 0.240 mmol) in methanol-methylene chloride (1:1, 6.0 mL) was added a 1.0 M aqueous methanesulfonic acid (0.48 mL, 0.48 mmol) and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-methylene chloride-diethyl ether to provide the title compound as a yellow crystalline powder (melting point: 247.0-249.0°C) (140.0 mg, yield: 76%).

<sup>1</sup>H-NMR [DMSO-d<sub>6</sub>, 120°C] (ammonium salt NH<sup>+</sup> protons were not observed) δ: 2.43(s, 6H), 3.05(s, 6H), 3.73(s, 4H), 3.76(s, 6H), 3.86(s, 12H), 7.16(s, 4H), 7.47(dd, J=3.1, 9.0Hz, 2H), 7.86(d, J=9.0Hz, 2H), 8.07(d, J=3.1Hz, 2H).

#### Example 8

N,N'-Bis[2-(4-ethoxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine dimethanesulfonate





Following the procedure of Example 1, N,N'-bis[2-(4-ethoxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine was obtained as a colorless oil (133.0 mg, yield: 79%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (150.0 mg, 0.280 mmol) synthesized in Reference Example 1 and 4-ethoxy-3,5-dimethoxyphenylboronic acid (140.0 mg, 0.620 mmol) synthesized in Reference Example 3.

To a solution of N,N'-bis[2-(4-ethoxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (133.0 mg, 0.220 mmol) in methanol (2.0 mL) was added a 1.0 M aqueous methanesulfonic acid solution (0.44 mL, 0.44 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to afford the title compound as a yellow crystalline powder (melting point: 232.0-234.0°C) (120.0 mg, yield: 54%).


<sup>1</sup>H-NMR [CDCl<sub>3</sub>] (ammonium salt NH<sup>+</sup> protons were not observed)

δ: 1.38(t, J=7.1Hz, 6H), 2.87(s, 6H), 3.32(s, 6H),  
3.80(s, 4H), 3.95(s, 12H), 4.12(q, J=7.1Hz, 4H),  
7.05(s, 4H), 7.70(dd, J=2.9, 9.3Hz, 2H),  
7.88(d, J=9.3, Hz, 2H), 8.71(d, J=2.9Hz, 2H).

#### Example 9

N,N'-Bis[2-(3,5-dimethoxy-4-propoxyphenyl)-5-pyridyl]-N,N'-

dimethylethylenediamine dimethanesulfonate



Following the procedure of Example 1, N,N'-bis[2-(3,5-dimethoxy-4-propoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine was obtained as a colorless oil (124.0 mg, yield: 98%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (108.0 mg, 0.200 mmol) synthesized in Reference Example 1 and 4-propoxy-3,5-dimethoxyphenylboronic acid (105.0 mg, 0.440 mmol) synthesized by a similar procedure as described in Reference Example 3.

To a solution of N,N'-bis[2-(3,5-dimethoxy-4-propoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (124.0 mg, 0.196 mmol) in methanol-chloroform (1:2, 3.0 mL) was added a 1.0 M aqueous methanesulfonic acid (0.41 mL, 0.41 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from ethanol-diethyl ether-hexane to afford the title compound as yellow a crystalline powder (melting point: 241.0-243.0°C ) (91.0 mg, yield: 56%).

<sup>1</sup>H-NMR [DMSO-d<sub>6</sub>, 120°C ] (ammonium salt NH<sup>+</sup> protons were not

observed) δ: 0.98(t, J=7.0Hz, 6H),


1.67(tq, J=7.0, 7.0Hz, 4H), 2.44(s, 6H), 3.06(s, 6H), 3.73(s, 4H), 3.85(s, 12H), 3.91(t, J=7.0Hz, 4H),

7.16(s, 4H), 7.47(dd, J=2.9, 9.0Hz, 2H),

7.86(d, J=9.0Hz, 2H), 8.06(d, J=2.9Hz, 2H).

Example 10

N,N'-Bis[2-(4-isopropoxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine dimethanesulfonate



Following the procedure of Example 1, N,N'-bis[2-(4-isopropoxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine was obtained as a colorless oil (74.0 mg, yield: 92%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (69.0 mg, 0.130 mmol) synthesized as described in Reference Example 1 and 4-isopropoxy-3,5-dimethoxyphenylboronic acid (61.0 mg, 0.260 mmol) synthesized by a similar procedure as described in Reference Example 3.

To a solution of N,N'-bis[2-(4-isopropoxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (74.0 mg, 0.12 mmol) in methanol (2.0 mL) was added a 1.0 M aqueous methanesulfonic acid solution (0.26 mL, 0.26 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to afford the title compound as a yellow crystalline powder (melting point: 250.0-253.0°C)(60.0 mg, yield: 57%).

<sup>1</sup>H-NMR [CDCl<sub>3</sub>](ammonium salt NH<sup>+</sup> protons were not observed)


δ: 1.31(d, J=6.1Hz, 12H), 2.86(s, 6H), 3.32(s, 6H),

3.80(s, 4H), 3.94(s, 12H), 4.46(qq, J=6.1, 6.1Hz, 2H), 7.06(s, 4H), 7.68(dd, J=3.2, 9.3Hz, 2H),

7.87(d, J=9.3Hz, 2H), 8.74(d, J=3.2Hz, 2H).

Example 11

N,N'-Bis[2-[3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)-phenyl]-5-pyridyl]-N,N'-dimethylethylenediamine dimethanesulfonate




Following the procedure of Example 1, N,N'-bis[2-[3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)phenyl]-5-pyridyl]-N,N'-dimethylethylenediamine was obtained as a colorless oil (80.0 mg, yield: 46%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (128.0 mg, 0.240 mmol) synthesized in Reference Example 1 and 3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)phenylboronic acid (146.0 mg, 0.530 mmol) synthesized as described in Reference Example 4.

To a solution of N,N'-bis[2-[3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)phenyl]-5-pyridyl]-N,N'-dimethylethylenediamine (80.0 mg, 0.11 mmol) in methanol (2.0 mL) was added a 1.0 M aqueous methanesulfonic acid (0.22 mL, 0.22 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to afford the title compound as a yellow crystalline powder (melting point: 253.0-257.0°C)(47.0 mg, yield: 22%).

$^1\text{H-NMR}$  [DMSO, 120°C] (ammonium salt  $\text{NH}^+$  protons were not observed)  $\delta$ : 2.41(s, 6H), 3.04(s, 6H), 3.70(s, 4H), 3.88(s, 12H), 4.46(q,  $^3J_{\text{HF}}=9.0\text{Hz}$ , 4H), 7.22(s, 4H), 7.32(dd,  $J=3.2, 9.0\text{Hz}$ , 2H), 7.80(d,  $J=9.0\text{Hz}$ , 2H), 8.11(d,  $J=3.2\text{Hz}$ , 2H).

Example 12

N,N'-Bis[2-[4-(2-methoxyethoxy)-3,5-dimethoxyphenyl]-5-pyridyl]-N,N'-dimethylethylenediamine dimethanesulfonate



Following the procedure of Example 1, N,N'-bis[2-[4-(2-methoxyethoxy)-3,5-dimethoxyphenyl]-5-pyridyl]-N,N'-dimethylethylenediamine was obtained as a colorless oil (139.0 mg, yield: 75%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (150.0 mg, 0.280 mmol) synthesized as described in Reference Example 1 and 4-(2-methoxyethoxy)-3,5-dimethoxyphenylboronic acid (158.0 mg, 0.620 mmol) synthesized as described in Reference Example 5.

To a solution of N,N'-bis[2-[4-(2-methoxyethoxy)-3,5-dimethoxyphenyl]-5-pyridyl]-N,N'-dimethylethylenediamine (139.0 mg, 0.21 mmol) in methanol (2.0 mL) was added a 1.0 M aqueous methanesulfonic acid solution (0.42 mL, 0.42 mmol) was added, and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to afford the title compound as a yellow crystalline powder (melting point: 211.0-214.0°C ) (127.0 mg, yield: 71%).

<sup>1</sup>H-NMR [CDCl<sub>3</sub>](ammonium salt NH<sup>+</sup> protons were not observed)

δ: 2.86(s, 6H), 3.32(s, 6H), 3.44(s, 6H),

3.69-3.74(m, 4H), 3.80(s, 4H), 3.95(s, 12H),


4.16-4.21(m, 4H), 7.05(s, 4H),

7.69(dd, J=2.9, 9.3Hz, 2H), 7.88(d, J=9.3Hz, 2H),

8.73(d, J=2.9Hz, 2H).

Example 13

N,N'-Bis[2-[4-(2-hydroxyethoxy)-3,5-dimethoxyphenyl]-5-pyridyl]-N,N'-  
dimethylethylenediamine



Following the procedure of Example 1, an oil (357.0 mg) containing N,N'-bis[2-[4-[2-(t-butyltrimethylsiloxy)ethoxy]-3,5-dimethoxyphenyl]-5-pyridyl]-N,N'-dimethylethylenediamine was obtained from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (162.0 mg, 0.300 mmol) synthesized as described in Reference Example 1 and 4-[2-(t-butyltrimethylsiloxy)ethoxy]-3,5-dimethoxyphenylboronic acid (374.0 mg, 1.05 mmol) synthesized as described in Reference Example 6.


To a solution of the oil (357.0 mg) obtained by the preceding procedure in acetonitrile (19 mL) was added 46% hydrofluoric acid (1.0 mL). After stirring for 3 hours, a 1.0 M aqueous sodium hydroxide was added to the reaction mixture, and the resulting mixture was concentrated under reduced pressure to remove the organic solvent. The residue was extracted with chloroform, and the organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to obtain an oil. Methanol-chloroform-diethyl ether-hexane was added to the oil and the resulting precipitate was collected by filtration to afford the title compound as a yellow crystalline powder (melting point: 178.0-180.5°C)[141.0 mg, yield: 74% based on N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-

dimethylethylenediamine].

<sup>1</sup>H-NMR [CDCl<sub>3</sub>] δ: 3.03(s, 6H), 3.53(br s, 2H), 3.67(s, 4H),  
3.69-3.79(m, 4H), 3.96(s, 12H), 4.14-4.25(m, 4H),  
6.99(dd, J=3.0, 8.8Hz, 2H), 7.18(s, 4H),  
7.55(d, J=8.8Hz, 2H), 8.21(d, J=3.0Hz, 2H).

#### Example 14

N,N'-Bis[2-(4-hydroxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-  
dimethylethylenediamine dimethanesulfonate



Following the procedure of Example 1, N,N'-bis[2-(4-benzyloxy-3,5-  
dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine was obtained as a colorless  
viscous oil (64.0 mg, yield: 63%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-  
N,N'-dimethylethylenediamine (75.0 mg, 0.140 mmol) synthesized as described in Reference  
Example 1 and 4-benzyloxy-3,5-dimethoxyphenylboronic acid (81.0 mg, 0.280 mmol).


To a solution of N,N'-bis[2-(4-benzyloxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-  
dimethylethylenediamine (64.0 mg, 0.090 mmol) in acetic acid (1.0 mL) was added 10%  
palladium on charcoal (43.0 mg). Under hydrogen, the mixture was stirred at 50°C for 1  
hour and 30 minutes. Insoluble materials were removed by filtration through Celite, and the  
filtrate was concentrated under reduced pressure. A solution of the residue in methanol-  
chloroform (1:10) was washed with saturated aqueous sodium hydrogencarbonate and brine,  
dried over anhydrous sodium sulfate, and concentrated under reduced pressure. To a

solution of the residue in methanol (2.0 mL) was added a 1.0 M aqueous methanesulfonic acid (0.20 mL, 0.20 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to afford the title compound as a yellow crystalline powder (melting point:  $\geq 300.0^\circ$ ) (20.0 mg, yield: 31% based on N,N'-bis[2-(4-benzyloxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine).

$^1\text{H-NMR}$  [DMSO,  $120^\circ\text{C}$ ] (neither ammonium salt  $\text{NH}^+$  protons nor phenol OH protons were observed)  $\delta$ : 2.38(s, 6H), 3.04(s, 6H), 3.70(s, 4H), 3.85(s, 12H), 7.15(s, 4H), 7.43(dd,  $J=2.9, 9.0\text{Hz}$ , 2H), 7.80(d,  $J=9.0\text{Hz}$ , 2H), 7.99(d,  $J=2.9\text{Hz}$ , 2H).

#### Example 15

N,N'-Bis[2-(4-butyryloxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine dimethanesulfonate



To a solution of N,N'-bis[2-(4-hydroxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (74.0 mg, 0.135 mmol) synthesized as described in Example 14 in pyridine (1.5 mL) was added butyric anhydride (101.0 mg, 0.640 mmol). After stirring at room temperature for 12 hours, the reaction mixture was concentrated under reduced pressure. Saturated aqueous sodium hydrogen carbonate was added to the residue, and the mixture was




extracted with chloroform. The organic layer was washed with brine, dried over anhydrous sodium carbonate, and concentrated under reduced pressure. To a solution of the residue in methanol (2.0 mL) was added a 1.0 M aqueous methanesulfonic acid (0.27 mL, 0.27 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from chloroform-diethyl ether to afford the title compound as a yellow crystalline powder (melting point: 196.0-200.0°C)(70.0 mg, yield: 59%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)(ammonium salt NH<sup>+</sup> protons were not observed)

δ: 1.07(t, J=7.3Hz, 6H), 1.82(tq, J=7.3, 7.3Hz, 4H),  
2.61(t, J=7.3Hz, 4 H), 2.85(s, 6H), 3.32(s, 6H),  
3.81(s, 4H), 3.92(s, 12H), 7.08(s, 4H),  
7.70(dd, J=2.9, 9.3Hz, 2H), 7.89(d, J=9.3Hz, 2H),  
8.72(d, J=2.9Hz, 2H).

#### Example 16

N,N'-Bis[2-(3-methoxy-4,5-methylenedioxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine dimethanesulfonate



Following the procedure of Example 1, N,N'-bis[2-(3-methoxy-4,5-methylenedioxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine was obtained as a colorless oil (109.0 mg, yield: 72%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (150.0 mg, 0.280 mmol) synthesized as described in


Reference Example 1 and 3-methoxy-4,5-methylenedioxyphenylboronic acid (115.0 mg, 0.590 mmol).

To a solution of N,N'-bis[2-(3-methoxy-4,5-methylene-dioxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (109.0 mg, 0.200 mmol) in methanol (3.0 mL) was added a 1.0 M aqueous solution of methanesulfonic acid (0.40 mL, 0.40 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to yield the title compound as a yellow crystalline powder (melting point: 275.0°-276.0°C)(109.0 mg, yield: 79%).

<sup>1</sup>H-NMR [CD<sub>3</sub>OD-CDCl<sub>3</sub>(1:10)](ammonium salt NH<sup>+</sup> protons were not observed) δ: 2.84(s, 6H), 3.20(s, 6H), 3.85(s, 4H), 4.02(s, 6H), 6.09(s, 4H), 6.93(d, J=1.7Hz, 2H), 7.06(d, J=1.7Hz, 2H), 7.74(dd, J=2.9, 9.5Hz, 2H), 7.82(d, J=9.5Hz, 2H), 8.34(d, J=2.9Hz, 2H).

#### Example 17

N,N'-Bis[2-(3,4,5-triethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine dimethanesulfonate



Following the procedure of Example 1, N,N'-bis[2-(3,4,5-triethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine was obtained as a pale yellow oil (123.0 mg, yield: 93%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-

dimethylethylenediamine (108.0 mg, 0.200 mmol) synthesized as described in Reference Example 1 and 3,4,5-triethoxyphenylboronic acid (122.0 mg, 0.480 mmol).

To a solution of N,N'-bis[2-(3,4,5-triethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (123.0 mg, 0.186 mmol) in methanol (3.0 mL) was added a 1.0 M solution of methanesulfonic acid in methanol (0.38 mL, 0.38 mmol), and the reaction mixture was concentrated under reduced pressure. Chloroform-diethyl ether was added to the residue and the resulting precipitate was collected by filtration to afford the title compound as a yellow crystalline powder [melting point: 237.0°C (decomposed)] (110.0 mg, yield: 70%).

<sup>1</sup>H-NMR [CDCl<sub>3</sub>] (ammonium salt NH<sup>+</sup> protons were not observed)

δ: 1.37(t, J=7.0Hz, 6H), 1.46(t, J=6.8Hz, 12H),

2.87(s, 6H), 3.31(s, 6H), 3.79(s, 4H),


4.12(q, J=7.0Hz, 4H), 4.17(q, J=6.8Hz, 8H),

7.01(s, 4H), 7.67(br d, J=9.0Hz, 2H),

7.82(d, J=9.0Hz, 2H), 8.69(br s, 2H).

#### Example 18

N,N'-Bis[2-(3,5-dimethoxy-4-methylthiophenyl)-5-pyridyl]-N,N'-dimethylethylenediamine dimethanesulfonate



Following the procedure of Example 1, N,N'-bis[2-(3,5-dimethoxy-4-methylthiophenyl)-5-pyridyl]-N,N'-dimethylethylenediamine was obtained as a colorless viscous oil (61.0 mg, yield: 80%) from N,N'-bis(2-trifluoromethane-sulfonyloxy-5-pyridyl)-

N,N'-dimethylethylenediamine (70.0 mg, 0.130 mmol) synthesized as described in Reference Example 1 and 3,5-dimethoxy-4-methylthiophenylboronic acid (60.0 mg, 0.260 mmol).

To a solution of N,N'-bis[2-(3,5-dimethoxy-4-methylthiophenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (61.0 mg, 0.100 mmol) in methanol (2.0 mL) was added a 1.0 M aqueous solution of methanesulfonic acid (0.20 mL, 0.20 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to yield the title compound as a yellow crystalline powder (melting point: 242.0-244.0°C)(41.0 mg, yield: 51%).


<sup>1</sup>H-NMR (CDCl<sub>3</sub>)(ammonium salt NH<sup>+</sup> protons were not observed)

δ: 2.42(s, 6H), 2.86(s, 6H), 3.34(s, 6H), 3.82(s, 4H),

4.00(s, 12H), 7.01(s, 4H), 7.72(dd, J=2.9, 9.3Hz, 2H), 7.93(d, J=9.3Hz, 2H), 8.79(d, J=2.9Hz, 2H).

#### Example 19

N,N'-Bis[2-[3,5-dimethoxy-4-(1-pyrrolidinyl)phenyl]-5-pyridyl]-N,N'-dimethylethylenediamine



Following the procedure as described in Example 1, an oil was obtained from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (120.0 mg, 0.220 mmol) synthesized as described in Reference Example 1 and (1-bromo-)3,5-dimethoxy-4-(1-pyrrolidinyl)benzene (138.0 mg, 0.550 mmol). Methanol was added to the

oil and the resulting precipitate was collected by filtration to yield the title compound as a pale brown crystalline powder (melting point: 221.0-225.0°C)(81.0 mg, yield: 56%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.89-1.97(m, 8H), 3.01(s, 6H),


3.28-3.37(m, 8H), 3.65(s, 4H), 3.91(s, 12H),

6.97(dd, J=2.9, 8.8Hz, 2H), 7.14(s, 4H),

7.54(d, J=8.8Hz, 2H), 8.21(d, J=2.9Hz, 2H).

#### Example 20

N,N'-Bis[2-(4-dimethylamino-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine



Following the procedure of Example 1, an oil was obtained from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (129.0 mg, 0.240 mmol) synthesized as described in Reference Example 1 and 4-dimethylamino-3,5-dimethoxyphenylboronic acid (135.0 mg, 0.600 mmol). Methanol was added to the oil and the resulting precipitate was collected by filtration to yield the title compound as a pale brown crystalline powder (melting point: 194.0-196.0°)(84.0 mg, yield: 58%).


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 2.85(s, 12H), 3.02(s, 6H), 3.66(s, 4H),

3.93(s, 12H), 6.98(dd, J=2.9, 9.0Hz, 2H), 7.12(s, 4H), 7.55(d, J=9.0Hz, 2H), 8.22(d,

J=2.9Hz, 2H).

#### Example 21

N,N'-Bis[2-(3,5-dimethoxy-4-methylphenyl)-5-pyridyl]-N,N'-  
dimethylethylenediamine dimethanesulfonate



Following the procedure of Example 7, N,N'-bis[2-(3,5-dimethoxy-4-methylphenyl)-  
5-pyridyl]-N,N'-dimethylethylenediamine was obtained as a colorless oil (37.0 mg, yield:  
40%) from 5-chloro-2-(3,5-dimethoxy-4-methylphenyl)pyridine (99.0 mg, 0.370 mmol)  
synthesized by a similar procedure as described in Reference Example 2.


To a solution of N,N'-bis[2-(3,5-dimethoxy-4-methylphenyl)-5-pyridyl]-N,N'-  
dimethylethylenediamine (37.0 mg, 0.070 mmol) in methanol (2.0 mL) was added a 1.0 M  
aqueous methanesulfonic acid (0.15 mL, 0.15 mmol), and the reaction mixture was  
concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the  
mixture was concentrated under reduced pressure. The residue was recrystallized from  
methanol-chloroform to yield the title compound as a yellow crystalline powder (melting  
point: 272.0-274.0°C)(22.0 mg, yield: 44%).

$^1\text{H-NMR}$  [ $\text{CD}_3\text{OD-CDCl}_3(1:10)$ ](ammonium salt  $\text{NH}^+$  protons were not  
observed)  $\delta$ : 2.12(s, 6H), 2.83(s, 6H), 3.24(s, 6H),  
3.85(s, 4H), 3.91(s, 12H), 7.23(s, 4H),  
7.76(dd,  $J=2.9, 9.3\text{Hz}$ , 2H), 7.94(d,  $J=9.3\text{Hz}$ , 2H),  
8.45(d,  $J=2.9\text{Hz}$ , 2H).

#### Example 22

N,N'-Bis[2-(4-fluoro-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-

dimethylethylenediamine dimethanesulfonate



Following the procedure of Example 1, an oil was obtained from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (150.0 mg, 0.280 mmol) synthesized as described in Reference Example 1 and 4-fluoro-3,5-dimethoxyphenylboronic acid (167.0 mg, 0.840 mmol). Hexane was added to the oil and the resulting precipitate was collected by filtration to provide N,N'-bis[2-(4-fluoro-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine as a yellow crystalline powder (160.0 mg; quantitative).

To a solution of N,N'-bis[2-(4-fluoro-3,5-dimethoxy-phenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (150.0 mg, 0.270 mmol) in chloroform (30 mL) was added a 1.0 M aqueous methanesulfonic acid solution (0.60 mL, 0.60 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. Diethyl ether was added to the residue and the resulting precipitate was collected by filtration to yield the title compound as a yellow crystalline powder (melting point:  $\geq 300.0^\circ\text{C}$ ) (181.0 mg, yield: 90%).

$^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ ) (ammonium salt  $\text{NH}^+$  protons were not observed)


$\delta$ : 2.33(s, 6H), 3.03(s, 6H), 3.77(br s, 4H),

3.89(s, 12H), 7.25(d,  $^4J_{\text{HF}}=7.0\text{Hz}$ , 4H),

7.50-7.59(m, 2H), 7.99-8.06(m, 4H).

Example 23

N,N'-Bis[2-(4-chloro-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-  
dimethylethylenediamine dimethanesulfonate



Following the procedure of Example 1, N,N'-bis[2-(4-chloro-3,5-dimethoxyphenyl)-  
5-pyridyl]-N,N'-dimethylethylenediamine was obtained as a pale yellow amorphous powder  
(101.0 mg, yield: 86%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-  
dimethylethylenediamine (108.0 mg, 0.200 mmol) synthesized as described in Reference  
Example 1 and 4-chloro-3,5-dimethoxyphenylboronic acid (95.0 mg, 0.440 mmol).


To a solution of N,N'-bis[2-(4-chloro-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-  
dimethylethylenediamine (78.0 mg, 0.130 mmol) in methanol-chloroform (1:2, 50 mL) was  
added a 1.0 M aqueous methanesulfonic acid solution (0.27 mL, 0.27 mmol), and the reaction  
mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the  
residue, and the mixture was concentrated under reduced pressure. The residue was  
recrystallized from methanol-chloroform-diethyl ether to yield the title compound was  
obtained as a yellow crystalline powder (melting point: 280.0-285.0°C) (83.0 mg, yield:  
80%).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 120°C) (ammonium salt NH<sup>+</sup> protons were not  
observed) δ: 2.45(s, 6H), 3.05(s, 6H), 3.72(s, 4H),  
3.91(s, 12H), 7.26(s, 4H), 7.37(dd, J=3.1, 9.0Hz, 2H), 7.87(d, J=9.0Hz, 2H), 8.14(d,  
J=3.1Hz, 2H).

Example 24



N,N'-Bis[2-(4-cyano-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-  
dimethylethylenediamine dimethanesulfonate




Following the procedure of Example 1, N,N'-bis[2-(4-cyano-3,5-dimethoxyphenyl)-  
5-pyridyl]-N,N'-dimethylethylenediamine was obtained as a colorless oil (125.0 mg, yield:  
79%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-  
dimethylethylenediamine (150.0 mg, 0.280 mmol) synthesized as described in Reference  
Example 1 and 4-cyano-3,5-dimethoxyphenylboronic acid (145.0 mg, 0.700 mmol).

To a solution of N,N'-bis[2-(4-cyano-3,5-dimethoxy-phenyl)-5-pyridyl]-N,N'-  
dimethylethylenediamine (125.0 mg, 0.220 mmol) in methanol (2.0 mL) was added a 1.0 M  
aqueous methanesulfonic acid solution (0.44 mL, 0.44 mmol), and the reaction mixture was  
concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the  
mixture was concentrated under reduced pressure. Methanol (2.5 mL) was added to the  
residue, and the resulting suspension was stirred in a bath maintained at 70°C. After  
cooling, the resulting precipitate was collected by filtration to afford the title compound as a  
yellow crystalline powder (melting point:  $\geq 300.0^\circ\text{C}$ )(72.0 mg, yield: 43%).

$^1\text{H-NMR}$  [ $\text{CD}_3\text{OD-CDCl}_3(1:10)$ ](ammonium salt  $\text{NH}^+$  protons were not  
observed)  $\delta$ : 2.35(s, 6H), 3.04(s, 6H), 3.77(s, 4H),  
3.97(s, 12H), 7.32(s, 4H), 7.28-7.34(m, 2H),  
8.06(d,  $J=9.0\text{Hz}$ , 2H), 8.19(d,  $J=3.2\text{Hz}$ , 2H).

Example 25

N,N'-Bis[2-[3,5-dimethoxy-4-(N-methoxy-N-methylcarbamoyl)phenyl]-5-pyridyl]-  
N,N'-dimethylethylenediamine




Following the procedure of Example 1, an oil was obtained from N,N'-bis(2-  
5 trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (76.0 mg, 0.140  
mmol) synthesized as described in Reference Example 1 and 3,5-dimethoxy-4-(N-methoxy-  
N-methylcarbamoyl)phenylboronic acid (91.0 mg, 0.340 mmol) synthesized in Reference  
Example 7. Chloroform-hexane was added to the oil and the resulting precipitate was  
collected by filtration to yield the title compound as a yellow crystalline powder (melting  
point: 217.0-220.0°C)(34.0 mg, yield: 36%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)(mixture of geometrical isomers of the amide,  
data for the major isomer) δ: 3.04(s, 6H),  
3.40(s, 6H), 3.51(s, 6H), 3.68(s, 4H), 3.91(s, 12H),  
7.01(dd, J=2.5, 8.6Hz, 2H), 7.15(s, 4H),  
15 7.59(d, J=8.6Hz, 2H), 8.23(d, J=2.5Hz, 2H).

#### Example 26

N,N'-Bis[2-(4-t-butoxycarbonyl-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-  
dimethylethylenediamine




Following the procedure of Example 1, crude crystals were obtained from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethylethylenediamine (167.0 mg, 0.300 mmol) synthesized as described in Reference Example 1 and 4-t-butoxy-carbonyl-3,5-dimethoxyphenylboronic acid (187.0 mg, 0.660 mmol). The crude crystals were  
5 recrystallized from methylene chloride-diethyl ether-hexane to yield the title compound as a colorless crystalline powder (melting point: 211.0-213.0°C)(195.0 mg, yield: 90%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.59(s, 18H), 3.02(s, 6H), 3.66(s, 4H),  
3.90(s, 12H), 6.98(dd, J=3.1, 8.7Hz, 2H), 7.11(s, 4H), 7.55(d, J=8.7Hz, 2H), 8.22(d, J=3.1Hz, 2H).

#### Example 27

N,N'-Bis[2-(4-carboxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-  
dimethylethylenediamine ditrifluoroacetate




To a solution of N,N'-bis[2-(4-t-butoxycarbonyl-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (72.0 mg, 0.100 mmol) synthesized as described in Example 26 in methylene chloride (1.0 mL) was added trifluoroacetic acid (1.0 mL), and the resulting mixture was stirred for 3 hours. Diethyl ether was added to the ice-cold reaction mixture and the resulting precipitate was collected by filtration to yield the title compound as a pale  
15 yellow crystalline powder [melting point: 235.0°C (decomposed)](72.0 mg, yield: 86%).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 120°C) (neither ammonium salt NH<sup>+</sup> protons nor carboxylic acid CO<sub>2</sub>H protons were observed) δ:

3.03(s, 6H), 3.68(s, 4H), 3.84(s, 12H),  
7.19(dd, J=3.1, 8.7Hz, 2H), 7.22(s, 4H),  
7.76(d, J=8.7Hz, 2H), 8.18(d, J=3.1Hz, 2H).

## 5 Example 28

N,N'-Bis[2-(4-ethoxycarboxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-  
dimethylethylenediamine




To a solution of N,N'-bis[2-(4-carboxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-  
dimethylethylenediamine ditrifluoroacetate (42.0 mg, 0.050 mmol) synthesized as described  
in Example 27 in methylene chloride (5.0 mL) were added dimethylformamide (0.1 mL, 1.29  
mmol) and oxalyl chloride (0.21 mL, 2.5 mmol). After the resulting mixture was stirred at  
room temperature for 30 minutes, ethanol (2.0 mL, 35 mmol) and N,N-diisopropyl-  
ethylamine (0.51 mL, 3.0 mmol) were added. The reaction mixture was stirred for 1 hour  
and concentrated under reduced pressure. The residue was purified by column  
chromatography on silica gel and crude crystals were obtained. The crude crystals were  
recrystallized from methylene chloride-diethyl ether-hexane to yield the title compound as a  
pale yellow crystalline powder (melting point: 206.0-208.0°C)(30.0 mg, yield: 86%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.38(t, J=7.0Hz, 6H), 3.03(s, 6H),  
3.67(s, 4H), 3.91(s, 12H), 4.40(q, J=7.0Hz, 4H),  
7.00(dd, J=2.9, 8.7Hz, 2H), 7.14(s, 4H),  
7.57(d, J=8.7Hz, 2H), 8.23(d, J=2.9Hz, 2H).

Example 29

N,N'-Bis[2-(4-acetyl-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-  
dimethylethylenediamine dimethanesulfonate



5 Under nitrogen, to a solution of (4-acetyl-3,5-dimethoxyphenyl)trimethyltin (235.0 mg, 0.760 mmol) synthesized as described in Reference Example 8 in toluene (10 mL) were added crude N,N'-bis(2-chloro-5-pyridyl)-N,N'-dimethylethylenediamine (78.0 mg, approx. 0.25 mmol) synthesized as described in Reference Example 9 and tetrakis(triphenylphosphine)palladium (116.0 mg, 0.100 mmol). After stirring at 130°C for 22 hours, a saturated aqueous potassium fluoride was added to the reaction mixture and the mixture was stirred for 1 hour. Insoluble materials were removed by filtration through Celite. The residue was washed with chloroform, and the aqueous layer of the filtrate was extracted with chloroform. The washing and the organic layer of the filtrate were combined, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield N,N'-bis[2-(4-acetyl-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine as a pale yellow crystalline powder [41.4 mg, yield: 23% based on N,N'-bis(2-hydroxy-5-pyridyl)-N,N'-dimethylethylenediamine ditrifluoroacetate].

To a solution of N,N'-bis[2-(4-acetyl-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethylethylenediamine (76.4 mg, 0.128 mmol) in chloroform (5.0 mL) was added a 1.0 M solution of methanesulfonic acid in methanol (0.32 mL, 0.32 mmol), and the reaction mixture was concentrated under reduced pressure. Chloroform-diethyl ether was added to the

residue and the resulting precipitate was collected by filtration to yield the title compound as a yellow crystalline powder (melting point: 261.0-265.5°C)(75.1 mg, yield: 74%).


<sup>1</sup>H-NMR (CDCl<sub>3</sub>)(ammonium salt NH<sup>+</sup> protons were not observed)

δ: 2.49(s, 6H), 2.86(s, 6H), 3.35(s, 6H), 3.80(s, 4H), 3.92(s, 12H), 7.03(s, 4H), 7.72(br d, J=9.2Hz, 2H),

7.94(d, J=9.2Hz, 2H), 8.82(br s, 2H).

### Example 30

1,4-Bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]hexahydro-1,4-diazepine  
dimethanesulfonate



Following the procedure of Example 7, 1,4-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]hexahydro-1,4-diazepine was obtained as a yellow amorphous powder (174.0 mg, yield: 59%) from 5-chloro-2-(3,4,5-trimethoxyphenyl)pyridine (335.0 mg, 1.20 mmol) synthesized as described in Reference Example 2 and hexahydro-1,4-diazepine (50.0 mg, 0.500 mmol).


To a solution of 1,4-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]hexahydro-1,4-diazepine (88.0 mg, 0.150 mmol) in methanol-methylene chloride (1:1, 4.0 mL) was added a 1.0 M aqueous methanesulfonic acid solution (0.30 mL, 0.30 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-methylene chloride-diethyl ether to yield the title compound as

yellow prisms (melting point: 246.0-248.0°C)(93.0 mg, yield: 79%).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 120°C) (ammonium salt NH<sup>+</sup> protons were not observed) δ: 2.04(tt, J=5.8, 5.8Hz, 2H), 2.42(s, 6H), 3.64(dd, J=5.8, 5.8Hz, 4H), 3.75(s, 6H), 3.83(s, 4H), 3.84(s, 12H), 7.14(s, 4H), 7.52(dd, J=3.1, 9.0Hz, 2H), 7.82(d, J=9.0Hz, 2H), 8.14(d, J=3.1Hz, 2H).

### Example 31

1,4-Bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]piperazine dimethanesulfonate



Following the procedure of Example 7, 1,4-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]piperazine was obtained as pale yellow needles (174.0 mg, yield: 61%) from 5-chloro-2-(3,4,5-trimethoxyphenyl)pyridine (335.0 mg, 1.20 mmol) synthesized as described in Reference Example 2 and piperazine (43.0 mg, 0.500 mmol).

To a solution of 1,4-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]piperazine (143.0 mg, 0.250 mmol) in methanol-methylene chloride (1:2, 6.0 mL) was added a 1.0 M aqueous methanesulfonic acid solution (0.50 mL, 0.50 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-chloroform-diethyl ether to yield the title compound as yellow needles [melting point: 262.0°C (decomposed)](171.0 mg, yield: 89%).


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) (data for the free base of the title compound)

δ: 3.46(s, 8H), 3.89(s, 6H), 3.96(s, 12H),

7.19(s, 4H), 7.32(dd, J=3.1, 8.7Hz, 2H),  
7.63(d, J=8.7Hz, 2H), 8.43(d, J=3.1Hz, 2H).

### Example 32

1,3-Bis[4-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-1-piperazinyl]propane  
tetrahydrochloride



Following the procedure of Example 7, 1,3-bis[4-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-1-piperazinyl]propane was obtained as a pale yellow oil (44.0 mg, yield: 25%) from 5-chloro-2-(3,4,5-trimethoxyphenyl)pyridine (166.0 mg, 0.600 mmol) synthesized in Reference Example 2 and 1,3-bis(1-piperazinyl)propane (90.0 mg, 0.250 mmol).

To a solution of 1,3-bis[4-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-1-piperazinyl]propane (44.0 mg, 0.060 mmol) in methanol-methylene chloride (1:2, 3.0 mL) was added 1.0 M hydrochloric acid solution (0.30 mL, 0.30 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to yield the title compound as a yellow crystalline powder [melting point: 264.0°C (decomposed)] (35.0 mg, yield: 69%).

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ ) (ammonium salt  $\text{NH}^+$  protons were not observed)


$\delta$ : 1.72-1.87(m, 2H), 2.46-2.56(m, 4H),  
2.61-2.71(m, 8H), 3.24-3.35(m, 8H), 3.88(s, 6H),  
3.95(s, 12H), 7.16(s, 4H), 7.24(dd, J=2.9, 8.7Hz, 2H),



7.58(d, J=8.7Hz, 2H), 8.37(d, J=2.9Hz, 2H).

### Example 33

N,N'-Bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]ethylenediamine




Following the procedure of Example 7, the title compound was obtained as a pale yellow amorphous powder (21.0 mg, yield: 7%) from 5-chloro-2-(3,4,5-trimethoxyphenyl)pyridine (335.0 mg, 1.20 mmol) synthesized as described in Reference Example 2 and ethylenediamine (30.0 mg, 0.500 mmol).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 3.50(s, 4H), 3.88(s, 6H), 3.95(s, 12H),  
4.01-4.09(br s, 2H), 7.01(dd, J=2.9, 8.5Hz, 2H),  
7.14(s, 4H), 7.53(d, J=8.5Hz, 2H), 8.16(d, J=2.9Hz, 2H).

### Example 34

cis-N,N'-Bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-1,2-cyclohexanediamine




Following the procedure of Example 7, the title compound was obtained as a pale yellow amorphous powder (37.0 mg, yield: 12%) from 5-chloro-2-(3,4,5-trimethoxyphenyl)pyridine (166.0 mg, 0.600 mmol) synthesized as described in Reference

Example 2 and cis-1,2-cyclohexanediamine (24.0 mg, 0.200 mmol).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.53-1.72(m, 6H), 1.79-1.91(m, 2H),  
3.75-3.85(m, 2H), 3.88(s, 6H), 3.94(s, 12H),  
3.94-4.02(m, 2H), 6.96(dd, J=2.6, 8.5Hz, 2H),  
7.13(s, 4H), 7.48(d, J=8.5Hz, 2H),  
8.13(d, J=2.6Hz, 2H).

### Example 35

trans-N,N'-Bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-1,4-cyclohexanediamine




Following the procedure of Example 7, crude crystals was obtained (66.0 mg, yield: 21%) from 5-chloro-2-(3,4,5-trimethoxyphenyl)pyridine (335.0 mg, 1.20 mmol) synthesized as described in Reference Example 2 and trans-1,4-cyclohexanediamine (57.0 mg, 0.500 mmol). The crude crystals were recrystallized from chloroform-diethyl ether-hexane to yield the title compound as a pale brown crystalline powder (melting point: 264.0-267.0°C)(27.0 mg, yield: 9%).

<sup>1</sup>H-NMR [CD<sub>3</sub>OD-CDCl<sub>3</sub>(1:10)] (amine NH protons were not observed) δ: 1.34-1.46(m, 4H), 2.19-2.31(m, 4H),  
3.27-3.35(m, 2H), 3.87(s, 6H), 3.95(s, 12H),  
7.00(dd, J=2.9, 8.7Hz, 2H), 7.07(s, 4H),  
7.51(d, J=8.7Hz, 2H), 8.03(d, J=2.9Hz, 2H).

Example 36

N,N'-Bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-diethylethylenediamine  
dimethanesulfonate




Following the procedure of Example 7, N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-diethylethylenediamine was obtained as a pale yellow crystalline powder (100.0 mg, yield: 33%) from 5-chloro-2-(3,4,5-trimethoxyphenyl)-pyridine (335.0 mg, 1.20 mmol) synthesized as described in Reference Example 2 and N,N'-diethylethylenediamine (58.0 mg, 0.500 mmol).

To a solution of N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-diethylethylenediamine (100.0 mg, 0.165 mmol) in methanol-chloroform (1:1, 6.0 mL) was added a 1.0 M aqueous methanesulfonic acid solution (0.33 mL, 0.33 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to yield the title compound as a yellow crystalline powder (melting point: 254.0-256.0°C)(90.0 mg, yield: 69%).

$^1\text{H-NMR}$  (DMSO- $d_6$ , 120°C)(ammonium salt  $\text{NH}^+$  protons were not observed)  $\delta$ : 1.17(t,  $J=6.8\text{Hz}$ , 6H), 2.41(s, 6H), 3.50(q,  $J=6.8\text{Hz}$ , 4H), 3.66(s, 4H), 3.76(s, 6H), 3.86(s, 12H), 7.18(s, 4H), 7.45(dd,  $J=2.9, 9.0\text{Hz}$ , 2H), 7.85(d,  $J=9.0\text{Hz}$ , 2H), 8.09(d,  $J=2.9\text{Hz}$ , 2H).

Example 37

N,N'-Bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine dimethanesulfonate




Following the procedure of Example 7, crude N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine was obtained as a pale brown oil (275.0 mg) from 5-chloro-2-(3,4,5-trimethoxyphenyl)pyridine (335.0 mg, 1.20 mmol) synthesized as described in Reference Example 2 and N,N'-dimethyl-1,3-propanediamine (65.0 mg, 0.500 mmol).

To a solution of the crude N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine (275.0 mg, approx. 0.46 mmol) in methanol-chloroform (1:1, 6.0 mL) was added a 1.0 M aqueous methanesulfonic acid solution (0.93 mL, 0.93 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to yield the title compound as yellow needles (melting point: 216.0-219.0°C)(255.0 mg, yield: 70% based on N,N'-dimethyl-1,3-propanediamine).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 120°C) (ammonium salt NH<sup>+</sup> protons were not observed) δ: 1.94(tt, J=7.3Hz, 2H), 2.43(s, 6H), 3.05(s, 6H), 3.52(t, J=7.3Hz, 4H), 3.76(s, 6H), 3.86(s, 12H), 7.15(s, 4H), 7.53(dd, J=3.1, 9.2Hz, 2H), 7.88(d, J=9.2Hz, 2H), 8.10(d, J=3.1Hz, 2H).

Example 38

N,N'-Bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-diethyl-1,3-propanediamine  
dimethanesulfonate




Following the procedure of Example 7, crude N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-diethyl-1,3-propanediamine was obtained as a pale brown crystalline powder (160.0 mg) from 5-chloro-2-(3,4,5-trimethoxyphenyl)pyridine (335.0 mg, 1.20 mmol) synthesized as described in Reference Example 2 and N,N'-diethyl-1,3-propanediamine (65.0 mg, 0.500 mmol).

To a solution of the crude N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-diethyl-1,3-propanediamine (160.0 mg, approx. 0.26 mmol) in methanol-chloroform (1:1, 6.0 mL) was added a 1.0 M aqueous methanesulfonic acid solution (0.51 mL, 0.51 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-methylene chloride-diethyl ether to yield the title compound as a yellow crystalline powder (melting point: 239.0-242.0°C)(92.0 mg, yield: 45% based on N,N'-diethyl-1,3-propanediamine).

$^1\text{H-NMR}$  ( $\text{DMSO-d}_6$ ,  $120^\circ\text{C}$ ) (ammonium salt  $\text{NH}^+$  protons were not observed)  $\delta$ : 1.14-1.21(m, 6H), 1.88-1.97(m, 2H), 2.43(s, 6H), 3.44-3.54(m, 8H), 3.76(s, 6H), 3.86(s, 12H), 7.16(s, 4H), 7.48-7.54(m, 2H), 7.83-7.90(m, 2H), 8.08-8.14(m, 2H).

Example 39

N,N'-Bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N',2,2-tetramethyl-1,3-propanediamine dimethanesulfonate




Following the procedure of Example 7, N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N',2,2-tetramethyl-1,3-propanediamine was obtained as a pale yellow oil (210.0 mg, yield: 68%) from 5-chloro-2-(3,4,5-trimethoxyphenyl)pyridine (335.0 mg, 1.20 mmol) synthesized as described in Reference Example 2 and N,N',2,2-tetramethyl-1,3-propanediamine (65.0 mg, 0.500 mmol).

To a solution of N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N',2,2-tetramethyl-1,3-propanediamine (210.0 mg, 0.340 mmol) in methanol-chloroform (1:1, 6.0 mL) was added a 1.0 M aqueous methanesulfonic acid (0.68 mL, 0.68 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to yield the title compound as a yellow crystalline powder (melting point: 123.0-128.0°C)(161.0 mg, yield: 59%).

$^1\text{H-NMR}$  ( $\text{DMSO-d}_6, 120^\circ\text{C}$ ) (ammonium salt  $\text{NH}^+$  protons were not observed)  $\delta$ : 1.08(s, 6H), 2.44(s, 6H), 3.15(s, 6H), 3.49(s, 4H), 3.76(s, 6H), 3.87(s, 12H), 7.18(s, 4H), 7.65(dd,  $J=3.1, 9.2\text{Hz}$ , 2H), 7.90(d,  $J=9.2\text{Hz}$ , 2H), 8.19(d,  $J=3.1\text{Hz}$ , 2H).

Example 40

2-Benzyloxy-N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine




5 Following the procedure of Example 7, the title compound was obtained as a pale yellow amorphous powder (69.0 mg, yield: 59%) from 5-chloro-2-(3,4,5-trimethoxyphenyl)pyridine (113.0 mg, 0.400 mmol) synthesized as described in Reference Example 2 and 2-benzyloxy-N,N'-dimethyl-1,3-propanediamine (35.0 mg, 0.160 mmol) synthesized as described in Reference Example 10.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 3.04(s, 6H), 3.49-3.51(m, 4H), 3.88(s, 6H), 3.93(s, 12H), 4.03-4.11(m, 1H), 4.52(s, 2H), 7.00(dd, J=3.1, 8.7Hz, 2H), 7.11-7.16(m, 2H), 7.16(s, 4H), 7.23-7.26(m, 3H), 7.49(d, J=8.7Hz, 2H), 8.21(d, J=3.1Hz, 2H).

15

Example 41

2-Hydroxy-N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine




To a solution of 2-benzyloxy-N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine (40.0 mg, 0.050 mmol) synthesized as described in Example 40 in methanol (5.0 mL) were added 1.0 M hydrochloric acid (2.5 mL, 2.5 mmol) and 10% palladium on charcoal (25.0 mg), and the resulting mixture was stirred at 45°C for 15 hours under hydrogen. Insoluble materials were removed by filtration through Celite. The filtrate was rendered basic by the addition of saturated aqueous sodium hydrogencarbonate and extracted with chloroform. The organic layer was washed with water, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield the title compound as a pale yellow amorphous powder (27.0 mg, yield: 89%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)(alcohol OH protons were not observed) δ:

3.08(s, 6H), 3.40-3.55(m, 4H), 3.88(s, 6H),  
3.94(s, 12H), 4.26-4.35(m, 1H),  
7.12(dd, J=2.9, 9.0Hz, 2H), 7.14(s, 4H),  
7.55(d, J=9.0Hz, 2H), 8.25(d, J=2.9Hz, 2H).

#### Example 42

~~N,N'-Bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N-methyl-2-aminoethyl]methylethylamine trihydrochloride~~



To a solution of methyliminodiacetic acid (74.0 mg, 0.503 mmol) in methylene chloride (2.5 mL) were added dimethylformamide (25 mg, 0.34 mmol) and oxalyl chloride



(0.22 mL, 2.3 mmol). The reaction mixture was stirred at room temperature for 2 hours and then concentrated under reduced pressure. To an ice-cold solution of the residue in methylene chloride (5 mL) were added N,N-diisopropylethylamine (0.90 mL, 5.2 mmol) and 5-amino-2-(3,4,5-trimethoxyphenyl)pyridine (262 mg, 1.00 mmol) synthesized as described in Reference Example 11. After being stirred in ice for 2 hours, saturated aqueous sodium hydrogencarbonate was added, and the mixture was extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield N,N-bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]carbamoylmethyl]methylamine as a colorless oil (94.5 mg, yield: 30%).

Under nitrogen, lithium aluminum hydride (23.1 mg, 0.609 mmol) was added to a solution of N,N-bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]carbamoylmethyl]methylamine (88.1 mg, 0.139 mmol) in anhydrous tetrahydrofuran (5.0 mL), and the resulting mixture was stirred at 65°C for 1 hour. Methanol (0.2 mL, 4.9 mmol) was added to the ice-cold reaction mixture, and the ice bath was removed. Water (0.2 mL), diethyl ether (20 mL), and anhydrous magnesium sulfate (1.5 g) were added, and the mixture was stirred at room temperature for 2 hours. Insoluble materials were removed by filtration through Celite, and the filtrate was concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield N,N-bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-2-aminoethyl]methylamine as a colorless oil (58.2 mg, yield: 69%).

To an ice-cold solution of N,N-bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-2-aminoethyl]methylamine (58.2 mg, 0.0964 mmol) in tetrahydrofuran (1.0 mL) were added paraformaldehyde (67.4 mg, 2.24 mmol) and sodium borohydride (46.4 mg, 1.23 mmol), followed by the gradual addition of trifluoroacetic acid (1.0 mL) over about 15 minutes.

The ice bath was removed, and the mixture was stirred at room temperature for 16 hours. Paraformaldehyde (33.2 mg, 1.11 mmol) and sodium borohydride (24.0 mg, 0.634 mmol) were added in several portions, and the resulting mixture was stirred at room temperature for additional 6 hours. To the ice-cold reaction mixture was added a 2.5 M aqueous sodium hydroxide (5.0 mL), and mixture was extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by preparative thin-layer chromatography on silica gel to yield N,N-bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N-methyl-2-aminoethyl]methylamine as a colorless oil (37.2 mg, yield: 61%).

To a solution of N,N-bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N-methyl-2-aminoethyl]methylamine (42.0 mg, 0.059 mmol) in ethanol (10 mL) was added 1.0 M hydrochloric acid (0.30 mL, 0.30 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (10 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from ethanol-diethyl ether to yield the title compound as a yellow crystalline powder [melting point: 187.0°C (decomposed)] (37.1 mg, yield: 75%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)(ammonium salt NH<sup>+</sup> protons were not observed)

δ: 3.07(br s, 3H), 3.19(br s, 6H), 3.49-3.70(m, 4H),

3.87(br s, 6H), 3.96(br s, 12H), 4.03-4.25(m, 4H),

7.17(br s, 4H), 7.83-7.97(m, 2H), 7.94-8.08(m, 2H),

8.16-8.30(m, 2H).


#### Example 43

~~N,N'-Bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N-methyl-3-~~

Ammon.  
a<sup>6</sup>

contd.  
a<sup>6</sup>

~~aminopropyl]methylanine~~




Following the procedure of Example 7, the title compound was obtained as a pale yellow amorphous powder (94.0 mg, yield: 28%) from 5-chloro-2-(3,4,5-trimethoxyphenyl)pyridine (335.0 mg, 1.20 mmol) synthesized as described in Reference Example 2 and N,N-bis(N-methyl-3-aminopropyl)methylamine (87.0 mg, 0.500 mmol).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.75-1.85(m, 4H), 2.23(s, 3H), 2.35-2.45(m, 4H), 2.99(s, 6H), 3.39-3.48(m, 4H), 3.87(s, 6H), 3.93(s, 12H), 7.04(dd, J=2.9, 8.7Hz, 2H), 7.14(s, 4H), 7.54(d, J=8.7Hz, 2H), 8.20(d, J=2.9Hz, 2H).

#### Example 44

Bis[N-2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-2-aminoethyl ether dimethanesulfonate



To an ice-cold solution of 5-amino-2-(3,4,5-trimethoxyphenyl)pyridine (203.6 mg, 0.783 mmol) synthesized as described in Reference Example 11 in methylene chloride (4 mL) were added N,N-diisopropylethylamine (0.20 mL, 1.1 mmol) and diglycolyl chloride (75.8 mg, 0.443 mmol). After being stirred in ice for 30 minutes, water was added to the reaction mixture. The mixture was extracted with methanol-chloroform (1:20), and the organic layer was dried over anhydrous sodium sulfate and concentrated under reduced

pressure. The residue was purified by column chromatography on silica gel to yield bis[N-  
[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]carbamoylmethyl]ether as a colorless oil (224.3 mg,  
yield: 93%).

Under nitrogen, lithium aluminum hydride (33.2 mg, 0.875 mmol) was added to a  
5 solution of bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]carbamoylmethyl]ether (121.3 mg,  
0.196 mmol) in anhydrous tetrahydrofuran (6 mL), and the mixture was stirred at 65°C for 2  
hours. Methanol (0.2 mL, 4.9 mmol) was added to the ice-cold reaction mixture, and the ice  
bath was removed. Water (0.2 mL), diethyl ether (20 mL), and anhydrous magnesium  
sulfate (1.5 g) were added, and the mixture was stirred at room temperature for 1 hour.  
Insoluble materials were removed by filtration through Celite, and the filtrate was  
concentrated under reduced pressure. The residue was purified by column chromatography  
on silica gel to yield bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-2-aminoethyl]ether as a  
colorless oil (98.5 mg, yield: 85%).

To a solution of bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-2-aminoethyl]ether  
(53.1 mg, 0.0900 mmol) in ethanol (5.0 mL) was added a 1.0 M aqueous methanesulfonic  
acid solution (0.20 mL, 0.20 mmol), and the reaction mixture was concentrated under  
reduced pressure. Ethanol (10 mL) was added to the residue, and the mixture was  
concentrated under reduced pressure. The residue was suspended in ethanol-diethyl ether  
and the resulting precipitate was collected by filtration to provide the title compound as a  
20 pale brown amorphous powder (49.9 mg, yield: 71%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)(ammonium salt NH<sup>+</sup> protons were not observed)

δ: 2.90(s, 6H), 3.38(br t, J=4.4Hz, 4H),


3.77(br t, J=4.4Hz, 4H), 3.89(s, 6H), 3.93(s, 12H),

6.93(s, 4H), 7.44(d, J=9.1Hz, 2H),

7.73(br d, J=9.1Hz, 2H), 8.38(br s, 2H).

#### Example 45

Bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N-methyl-2-aminoethyl]ether  
dimethanesulfonate



To an ice-cold solution of bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-2-aminoethyl]ether (23.6 mg, 0.0400 mmol) synthesized as described in Example 44 in tetrahydrofuran (0.5 mL) were added paraformaldehyde (25.9 mg, 0.862 mmol) and sodium borohydride (18.0 mg, 0.476 mmol). After a solution of trifluoroacetic acid (0.25 mL) in tetrahydrofuran (0.50 mL) was gradually added over about 10 minutes, the ice bath was removed and the mixture was stirred at room temperature for 30 minutes. Paraformaldehyde (25.0 mg, 0.833 mmol) and sodium borohydride (17.8 mg, 0.47 mmol) were added in several portions, and the resulting mixture was stirred for 20 hours. To the ice-cold reaction mixture was added a 2.5 M aqueous sodium hydroxide solution (5.0 mL), and the mixture was extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was purified by preparative thin-layer chromatography on silica gel to yield bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N-methyl-2-aminoethyl]ether as a colorless oil (18.8 mg, yield: 76%).

To a solution of bis[N-[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N-methyl-2-aminoethyl]ether (47.5 mg, 0.0769 mmol) in ethanol (5.0 mL) was added a 1.0 M aqueous


methanesulfonic acid solution (0.17 mL, 0.17 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (10 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from ethanol-diethyl ether to provide the title compound as a yellow crystalline powder (melting point: 165.5-169.0°C) (38.8 mg, yield: 62%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)(ammonium salt NH<sup>+</sup> protons were not observed)

δ: 2.89(s, 6H), 3.07(s, 6H), 3.63(br t, J=4.9Hz, 4H), 3.77(br t, J=4.9Hz, 4H), 3.88(s, 6H), 3.94(s, 12H), 6.98(s, 4H), 7.74(br dd, J=2.9, 9.3Hz, 2H), 7.87(br d, J=9.3Hz, 2H), 8.40(br s, 2H).

#### Example 46

N,N'-Bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-3-(1-pyrrolidinyl)-1,5-pentanediamine trihydrochloride



Following the procedure of Example 7, N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-3-(1-pyrrolidinyl)-1,5-pentanediamine was obtained as a colorless oil (49.6 mg, yield: 27%) from 2-(3,4,5-trimethoxyphenyl)-5-chloropyridine (182.6 mg, 0.635 mmol) synthesized as described in Reference Example 2 and N,N'-dimethyl-3-(1-pyrrolidinyl)-1,5-pentanediamine (54.0 mg, 0.217 mmol).


To a solution of N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-3-(1-pyrrolidinyl)-1,5-pentanediamine (49.6 mg, 0.0724 mmol) in ethanol (5.0 mL) was added

1.0 M hydrochloric acid solution (0.40 mL, 0.40 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (10 mL) was added to the residue, and the resulting mixture was concentrated under reduced pressure to yield the title compound as a pale yellow amorphous powder (46.0 mg, yield: 71%).

5 <sup>1</sup>H-NMR (CDCl<sub>3</sub>)(ammonium salt NH<sup>+</sup> protons were not observed)  
δ: 2.00-2.35(m, 8H), 3.10-3.25(m, 2H), 3.17(br s, 6H), 3.65-3.80(m, 3H),  
3.82-3.98(m, 4H), 3.90(s, 6H), 3.99(s, 12H), 7.24(s, 4H), 7.79(br d, J=8.8Hz, 2H),  
7.90(br d, J=8.8Hz, 2H), 8.20(br s, 2H).

#### Example 47

2-(Dimethylamino)methyl-1,4-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]piperazine trihydrochloride



Following the procedure of Example 7, 2-(dimethylamino)methyl-1,4-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]piperazine was obtained as a colorless oil (32.3 mg, yield: 20%)  
15 from 2-(3,4,5-trimethoxyphenyl)-5-chloropyridine (198.9 mg, 0.712 mmol) synthesized as described in Reference Example 2 and 2-(dimethylaminomethyl)piperazine (37.0 mg, 0.259 mmol).


To a solution of 2-(dimethylamino)methyl-1,4-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]piperazine (32.3 mg, 0.0512 mmol) in ethanol (5.0 mL) was added 1.0 M  
20 hydrochloric acid (0.20 mL, 0.20 mmol), and the reaction mixture was concentrated under

reduced pressure. Ethanol (10 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from ethanol-diethyl ether to yield the title compound as a pale yellow crystalline powder [melting point: 199.5°C (decomposed)](25.7 mg, yield: 68%).

5 <sup>1</sup>H-NMR (CDCl<sub>3</sub>)(data for the free base of the title compound)  
δ: 2.21(br d, J=11.5Hz, 1H), 2.24-2.33(m, 1H), 2.27(s, 6H),  
2.95(dd, J=11.5, 11.5Hz, 1H), 3.14(ddd, J=3.5, 11.5, 11.5Hz, 1H),  
3.18(dd, J=3.5, 11.5Hz, 1H), 3.34(ddd, J=3.5, 11.5, 11.5Hz, 1H),  
3.57(br ddd, J=3.5, 3.5, 11.5Hz, 1H), 3.74(br d, J=11.5 Hz, 1H),  
3.90(s, 6H), 3.97(s, 12H), 4.11(br d, J=11.5Hz, 1H), 7.19(s, 2H),  
7.19(s, 2H), 7.26(dd, J=2.9, 8.8Hz, 1H), 7.33(dd, J=2.9, 8.8Hz, 1H),  
7.61(d, J=8.8Hz, 1H), 7.62(d, J=8.8Hz, 1H),  
8.38(d, J=2.9Hz, 1H), 8.43(d, J=2.9Hz, 1H).

#### Example 48

N,N'-Bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-4,4'-bipiperidine dimethanesulfonate



Following the procedure of Example 7, N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-4,4'-bipiperidine was obtained as a pale yellow crystalline powder (228.0 mg, yield: 69%) from 5-chloro-2-(3,4,5-trimethoxyphenyl)pyridine (335.0 mg, 1.20 mmol) synthesized as described in Reference Example 2 and 4,4'-bipiperidine (84.0 mg, 0.500 mmol).

To a solution of N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-4,4'-bipiperidine




(228.0 mg, 0.340 mmol) in methanol-chloroform(1:1, 6.0 mL) was added a 1.0 M aqueous methanesulfonic acid (0.70 mL, 0.70 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-methylene chloride-diethyl ether to yield the title compound as yellow needles (melting point: 226.0-228.0°C)(170.0 mg, yield: 59%).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 120°C )(ammonium salt NH<sup>+</sup> protons were not observed) δ: 1.35-1.50(m, 6H), 1.81-1.92(m, 4H), 2.44(s, 6H), 2.81-2.97(m, 4H), 3.77(s, 6H), 3.88(s, 12H), 3.90-3.92(m, 4H), 7.20(s, 4H), 7.73(dd, J=2.9, 9.0Hz, 2H), 7.93(d, J=9.0Hz, 2H), 8.25(d, J=2.9Hz, 2H).

#### Example 49

N,N'-Bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,6-hexanediamine dimethanesulfonate




Following the procedure of Example 7, N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,6-hexanediamine was obtained as a pale yellow crystalline powder (243.0 mg, yield: 75%) from 5-chloro-2-(3,4,5-trimethoxyphenyl)pyridine (335.0 mg, 1.20 mmol) synthesized as described in Reference Example 2 and N,N'-dimethyl-1,6-hexanediamine (72.0 mg, 0.500 mmol).

To a solution of N,N'-bis[2-(3,4,5-trimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,6-hexanediamine (243.0 mg, 0.370 mmol) in methanol-methylene chloride (6:1, 3.5 mL) was added a 1.0 M aqueous methanesulfonic acid (0.76 mL, 0.76 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to yield the title compound as yellow needles (melting point: 192.0-195.0°C)(251.0 mg, yield: 81%).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 120°C) (ammonium salt NH<sup>+</sup> protons were not observed) δ: 1.37-1.45(m, 4H), 1.56-1.67(m, 4H), 2.42(s, 6H), 3.02(s, 6H), 3.40-3.47(m, 4H), 3.76(s, 6H), 3.87(s, 12H), 7.17(s, 4H), 7.50(dd, J=3.1, 9.0Hz, 2H), 7.89(d, J=9.0Hz, 2H), 8.04(d, J=3.1Hz, 2H).

#### Example 50

N,N'-Bis[2-(3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-diethylethylenediamine dimethanesulfonate



Following the procedure of Example 1, N,N'-bis[2-(3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-diethylethylenediamine was obtained as a colorless oil (55.0 mg, yield: 42%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-diethylethylenediamine (136.0 mg, 0.240 mmol) synthesized by a similar procedure to that described in Reference


Example 1 and 3,5-dimethoxyphenylboronic acid (80.0 mg, 0.530 mmol).

To a solution of N,N'-bis[2-(3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-diethylethylenediamine (55.0 mg, 0.100 mmol) in methanol (2.0 mL) was added a 1.0 M aqueous methanesulfonic acid (0.20 mL, 0.20 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to yield the title compound as a yellow crystalline powder (melting point: 280.0-282.0°C) (48.0 mg, yield: 64%).

<sup>1</sup>H-NMR [CD<sub>3</sub>OD-CDCl<sub>3</sub>, (1:10)] (ammonium salt NH<sup>+</sup> protons were not observed) δ: 1.26(t, J=6.8Hz, 6H), 2.83(s, 6H), 3.64(q, J=6.8Hz, 4H), 3.77(s, 4H), 3.89(s, 12H), 6.60(t, J=2.2Hz, 2H), 6.93(d, J=2.2Hz, 4H), 7.75(dd, J=2.9, 10.3Hz, 2H), 7.92(d, J=10.3Hz, 2H), 8.45(d, J=2.9Hz, 2H).

#### Example 51

N,N'-Diethyl-N,N'-bis[2-(4-isopropoxy-3,5-dimethoxyphenyl)-5-pyridyl]ethylenediamine dimethanesulfonate



Following the procedure of Example 1, N,N'-diethyl-N,N'-bis[2-(4-isopropoxy-3,5-dimethoxyphenyl)-5-pyridyl]ethylenediamine was obtained as a colorless oil (129.0 mg, yield: 98%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-

diethylethylenediamine (113.0 mg, 0.200 mmol) synthesized by a similar procedure as described in Reference Example 1 and 4-isopropoxy-3,5-dimethoxyphenylboronic acid (106.0 mg, 0.440 mmol).

To a solution of N,N'-diethyl-N,N'-bis[2-(4-isopropoxy-3,5-dimethoxyphenyl)-5-pyridyl]ethylenediamine (129.0 mg, 0.200 mmol) in methanol (2.0 mL) was added a 1.0 M aqueous methanesulfonic acid solution (0.40 mL, 0.40 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to afford the title compound as a yellow crystalline powder (melting point: 260.0-261.0°C) (102.0 mg, yield: 60%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)(ammonium salt NH<sup>+</sup> protons were not observed)

δ: 1.27(t, J=6.8Hz, 6H), 1.32(d, J=6.1Hz, 12H),

2.85(s, 6H), 3.72(s, 4H), 3.76(q, J=6.8Hz, 4H),


3.94(s, 12H), 4.47(qq, J=6.1, 6.1Hz, 2H), 7.07(s, 4H), 7.69(dd, J=2.9, 9.3Hz, 2H),

7.88(d, J=9.3Hz, 2H),

8.72(d, J=2.9Hz, 2H).

#### Example 52

N,N'-Bis[2-(3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine dimethanesulfonate



Following the procedure of Example 1, N,N'-bis[2-(3,5-dimethoxyphenyl)-5-

pyridyl]-N,N'-dimethyl-1,3-propanediamine was obtained as a colorless oil (35.0 mg, yield: 35%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethyl-1,3-propanediamine (95.0 mg, 0.190 mmol) synthesized by a similar procedure to that described in Reference Example 1 and 3,5-dimethoxyphenylboronic acid (63.0 mg, 0.420 mmol).

5 To a solution of N,N'-bis[2-(3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine (35.0 mg, 0.070 mmol) in methanol (2.0 mL) was added a 1.0 M aqueous methanesulfonic acid (0.15 mL, 0.15 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from ethanol to afford the title compound as a yellow crystalline powder (melting point: 228.0-231.0°C) (21.0 mg, yield: 43%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)(ammonium salt NH<sup>+</sup> protons were not observed)

δ: 2.07(tt, J=7.3, 7.3Hz, 2H), 2.90(s, 6H),

3.21(s, 6H), 3.58(t, J=7.3Hz, 4H), 3.85(s, 12H),


6.78(t, J=2.2Hz, 2H), 6.86(d, J=2.2Hz, 4H),

7.81(dd, J=3.2, 9.3Hz, 2H), 7.91(d, J=9.3Hz, 2H),

8.71(d, J=3.2Hz, 2H).

### Example 53

20 N,N'-Bis[2-(4-isopropoxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine dimethanesulfonate




Following the procedure of Example 1, N,N'-bis[2-(4-isopropoxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine was obtained as a colorless viscous oil (27.0 mg, yield: 33%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethyl-1,3-propanediamine (61.0 mg, 0.125 mmol) synthesized by a similar  
5 procedure to that described in Reference Example 1 and 4-isopropoxy-3,5-dimethoxyphenylboronic acid (66.0 mg, 0.275 mmol).

To a solution of N,N'-bis[2-(4-isopropoxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine (27.0 mg, 0.040 mmol) in methanol (2.0 mL) was added a 1.0 M aqueous methanesulfonic acid (0.10 mL, 0.10 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was recrystallized from methanol-diethyl ether to afford the title compound as a yellow crystalline powder (melting point: 191.0-192.0°C)(15.0 mg, yield: 42%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)(data for the free base of the title compound) δ: 1.31(d, J=6.1Hz, 12H), 1.96(tt, J=7.6, 7.6Hz, 2H), 3.01(s, 6H), 3.45(t, J=7.6 Hz, 4H), 3.91(s, 12H), 4.40(qq, J=6.1Hz, 2H), 7.14(s, 4H), 7.48(dd, J=2.9, 7.8Hz, 2H), 7.54(d, J=7.8Hz, 2H), 8.20(d, J=2.9Hz, 2H).

#### 20 Example 54

N,N'-Bis[2-(4-hydroxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine dimethanesulfonate



Following the procedure of Example 1, N,N'-bis[2-[4-(4-methoxybenzyloxy)-3,5-dimethoxyphenyl]-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine was obtained as a pale brown oil (75.0 mg, yield: 23%) from N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-N,N'-dimethyl-1,3-propanediamine (220.0 mg, 0.400 mmol) synthesized by a similar procedure to that described in Reference Example 1 and 4-(4-methoxybenzyloxy)-3,5-dimethoxyphenylboronic acid (308.0 mg, 0.970 mmol) synthesized by a similar procedure to that described in Reference Example 3.

To a solution of N,N'-bis[2-[4-(4-methoxybenzyloxy)-3,5-dimethoxyphenyl]-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine (72.0 mg, 0.089 mmol) in methylene chloride (1.0 mL) was added trifluoroacetic acid (1.0 mL). The reaction mixture was stirred at room temperature for 30 minutes, and concentrated under reduced pressure. A solution of the residue in chloroform was washed with saturated aqueous sodium hydrogencarbonate and brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield N,N'-bis[2-(4-hydroxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine as a pale yellow oil (43.0 mg, yield: 86%).

To a solution of N,N'-bis[2-(4-hydroxy-3,5-dimethoxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine (43.0 mg, 0.076 mmol) in methanol-chloroform (1:2, 3.0 mL) was added a 1.0 M aqueous methanesulfonic acid (0.16 mL, 0.16 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was


recrystallized from ethanol-diethyl ether to provide the title compound as a yellow crystalline powder (melting point: 198.0-202.0°C)(17.0 mg, yield: 30%).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 120°C )(neither ammonium salt NH<sup>+</sup> protons nor phenol OH protons were observed) δ:

1.93(tt, J=7.0, 7.0Hz, 2H), 2.41(s, 6H), 3.05(s, 6H), 3.52(t, J=7.0Hz, 4H), 3.85(s, 12H), 7.15(s, 4H), 7.59(dd, J=2.9, 9.2Hz, 2H), 7.88(d, J=9.2Hz, 2H), 8.09(d, J=2.9Hz, 2H).

#### Example 55

N,N'-Bis[2-(3-methoxy-4,5-methylenedioxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine dimethanesulfonate



Following the procedure of Example 1, N,N'-bis[2-(3-methoxy-4,5-methylenedioxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine was obtained as a colorless oil (46.0 mg, yield: 52%) from N,N'-dimethyl-N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-1,3-propanediamine (78.0 mg, 0.160 mmol) synthesized by a similar procedure to that described in Reference Example 1 and 3-methoxy-4,5-methylenedioxyphenylboronic acid (66.0 mg, 0.340 mmol).

To a solution of N,N'-bis[2-(3-methoxy-4,5-methylenedioxyphenyl)-5-pyridyl]-N,N'-dimethyl-1,3-propanediamine (46.0 mg, 0.080 mmol) in methanol (2.0 mL) was added a 1.0 M aqueous methanesulfonic acid solution (0.20 mL, 0.20 mmol), and the reaction mixture




was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. Diethyl ether was added to a solution of the residue in methanol and the resulting precipitate was collected by filtration to yield the title compound as a yellow amorphous powder (29.0 mg, yield: 46%).

5 <sup>1</sup>H-NMR [CD<sub>3</sub>OD-CDCl<sub>3</sub>(1:10)](ammonium salt NH<sup>+</sup> protons were not observed) δ: 2.04(tt, J=7.3, 7.3Hz, 2H), 2.86(s, 6H), 3.17(s, 6H), 3.58(t, J=7.3Hz, 4H), 3.99(s, 6H), 6.08(s, 4H), 6.87(d, J=1.7Hz, 2H), 6.97(d, J=1.7Hz, 2H), 7.78(dd, J=3.2, 9.5Hz, 2H), 7.88(d, J=9.5Hz, 2H), 8.38(d, J=3.2Hz, 2H).

#### Example 56

N,N'-Diethyl-N,N'-bis[2-(4-isopropoxy-3,5-dimethoxyphenyl)-5-pyridyl]-1,3-propanediamine dimethanesulfonate



15 Following the procedure of Example 1, N,N'-diethyl-N,N'-bis[2-(4-isopropoxy-3,5-dimethoxyphenyl)-5-pyridyl]-1,3-propanediamine was obtained as a pale yellow oil (78.0 mg, yield: 58%) from N,N'-diethyl-N,N'-bis(2-trifluoromethanesulfonyloxy-5-pyridyl)-1,3-propanediamine (116.0 mg, 0.200 mmol) synthesized by a similar procedure to that described in Reference Example 1 and 4-isopropoxy-3,5-dimethoxyphenylboronic acid (96.0 mg, 0.400 mmol).

To a solution of N,N'-diethyl-N,N'-bis[2-(4-isopropoxy-3,5-dimethoxyphenyl)-5-


pyridyl]-1,3-propanediamine (78.0 mg, 0.110 mmol) in methanol-chloroform (2:1, 3.0 mL) was added a 1.0 M aqueous methanesulfonic acid (0.24 mL, 0.24 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (5.0 mL) was added to the residue, and the mixture was concentrated under reduced pressure. The residue was

5 recrystallized from methanol-diethyl ether-hexane to yield the title compound as yellow flakes (melting point: 173.0-175.0°C) (66.0 mg, yield: 66%).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 120°C) (ammonium salt NH<sup>+</sup> protons were not observed) δ: 1.18(t, J=7.0Hz, 6H),  
 1.23(d, J=6.0Hz, 12H), 1.93(tt, J=7.0, 7.0Hz, 2H),  
 2.42(s, 6H), 3.47-3.56(m, 4H), 3.50(q, J=7.0Hz, 4H),  
 3.85(s, 12H), 4.38(qq, J=6.0, 6.0 Hz, 2H),  
 7.15(s, 4H), 7.56(dd, J=3.1, 9.0Hz, 2H),  
 7.90(d, J=9.0Hz, 2H), 8.09(d, J=3.1Hz, 2H).

#### Example 57

~~N,N'-Bis[N-[2-[3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)phenyl]-5-pyridyl]-N-methyl-3-aminopropyl]methanamine trihydrochloride~~



To a solution of 5-amino-2-[3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)phenyl]pyridine

20 (melting point: 176.0-177.5°C) (28.2 g, 85.4 mmol) synthesized by a similar procedure to that described in Reference Example 11 in methylene chloride (550 mL) was added N,N-diisopropylethylamine (18.0 mL, 103 mmol). To the ice-cold stirred solution was added

acryloyl chloride (7.00 mL, 86.0 mmol) dropwise over about 10 minutes. A saturated aqueous sodium hydrogencarbonate was added, and the resulting mixture was extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure. Chloroform-hexane (1:2, 300 mL) was added to the residue and the resulting precipitate was collected by filtration to yield crude 5-acryloylamino-2-[3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)phenyl]pyridine was obtained as colorless needles (melting point: 207.0-209.0°C)(38.2 g).

To a 30% solution of methylamine in ethanol (100 mL, 970 mmol) was added the crude 5-acryloylamino-2-[3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)phenyl]pyridine (19.2 g, approx. 50.2 mmol) obtained by the above-described procedure. After being stirred at room temperature for 1 hour, the reaction mixture was concentrated under reduced pressure. Chloroform (100 mL) was added to the residue, and the mixture was again concentrated under reduced pressure. To a solution of the residue in chloroform (100 mL) was added the crude 5-acryloylamino-2-[3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)phenyl]pyridine (16.0 g, approx. 41.8 mmol) obtained by the above-described procedure, and the mixture was stirred at 60°C for 12 hours. The solidified reaction mixture was dissolved in chloroform (600 mL), and the resulting solution was heated at 80°C for 2 hours, then the solvent was removed by distillation. The residue was again dissolved in chloroform (600 mL) and the solution was concentrated to about 200 mL. Hexane (100 mL) was added to the solution stirred at 80°C.

After cooling, the precipitated crystals were collected by filtration to yield N,N-bis[2-[N-[2-[3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)phenyl]-5-pyridyl]carbamoyl]ethyl]methylamine as a colorless crystalline powder [melting point: 99.0°C(decomposed)] [22.67 g, yield: 72% based on 5-amino-2-[3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)phenyl]pyridine].

Under nitrogen, lithium aluminum hydride (4.28 g, 113 mmol) was added to an ice-

10693530.0595001  
cold solution of N,N-bis[2-[N-[2-[3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)phenyl]-5-pyridyl]carbamoyl]ethyl]methylamine (22.50 g, 28.3 mmol) in anhydrous tetrahydrofuran (500 mL). The ice bath was removed, and the reaction mixture was stirred at 65°C for 1 hour. Methanol (20 mL, 490 mmol) was added to the ice-cold reaction mixture, and the ice bath was removed. Water (20 mL), diethyl ether (2 L), and anhydrous magnesium sulfate (130 g) were added, and the mixture was stirred overnight at room temperature. Insoluble materials were removed by filtration through Celite, and the filtrate was concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to provide N,N-bis[N-[2-[3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)phenyl]-5-pyridyl]-3-aminopropyl]methylamine as a colorless oil (20.32 mg, yield: 94%).

To a solution of N,N-bis[N-[2-[3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)phenyl]-5-pyridyl]-3-aminopropyl]methylamine (2.11 g, 2.75 mmol) in acetonitrile (10 mL) were added formalin (2.1 mL, 28 mmol) and sodium cyanoborohydride (0.55 g, 8.8 mmol). To the mixture was added to a solution of acetic acid (0.3 mL) in acetonitrile (1 mL) dropwise over about 10 minutes, and the resulting mixture was stirred at room temperature for 1 hour. To the reaction mixture were added formalin (2.1 mL, 28 mmol) and sodium cyanoborohydride (0.65 g, 10 mmol), then was added a solution of acetic acid (0.6 mL) in acetonitrile (0.5 mL) dropwise over about 10 minutes. After stirring at room temperature for 1 hour, 1.0 M aqueous sodium hydroxide solution (100 mL) was added to the reaction mixture, and the mixture was extracted with methylene chloride. The organic layer was concentrated under reduced pressure, and a solution of the residue in pyridine (15 mL) was stirred at 45°C for 1 hour, and then concentrated under reduced pressure. Toluene was added to the residue, and the mixture was concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to yield N,N-bis[N-[2-[3,5-dimethoxy-4-(2,2,2-

trifluoroethoxy)phenyl]-5-pyridyl]-N-methyl-3-aminopropyl]methylaniline was obtained as a colorless oil (984 mg, yield: 45%).

To a solution of N,N-bis[N-[2-[3,5-dimethoxy-4-(2,2,2-trifluoroethoxy)phenyl]-5-pyridyl]-N-methyl-3-aminopropyl]methylaniline (3.62 g, 4.55 mmol) in ethanol (50 mL) was added concentrated hydrochloric acid (1.4 mL, 17 mmol), and the reaction mixture was concentrated under reduced pressure. Ethanol (50 mL) was added to the residue, and the mixture was concentrated under reduced pressure. Ethyl acetate (20 mL) was added to the residue, followed by concentration of the resultant mixture under reduced pressure. The residue was suspended in ethyl acetate and collected by filtration to yield the title compound as a yellow crystalline powder [melting point: 140.0°C (decomposed)] (3.55 g, yield: 86%).

<sup>1</sup>H-NMR (CD<sub>3</sub>OD)(ammonium salt NH<sup>+</sup> protons were not observed)

δ: 2.17(br dddd, J=7.3, 7.3, 7.3, 7.3Hz, 4H),

2.94(s, 3H), 3.20(s, 6H), 3.23-3.32(m, 2H),

3.34-3.44(m, 2H), 3.65(dt, J=14.3, 7.3Hz, 2H),

3.70(dt, J=14.3, 7.3Hz, 2H), 3.97(s, 12H),

4.45(q, <sup>3</sup>J<sub>HF</sub>=8.8Hz, 4H), 7.15(s, 4H),


8.00(br dd, J=3.0, 9.5Hz, 2H),

8.12(br d, J=3.0Hz, 2H), 8.14(br d, J=9.5Hz, 2H).

## 20 Example 58

~~N,N'-Bis[N-[2-(3,5-dimethoxy-4-methylthiophenyl)-5-pyridyl]-N-methyl-3-aminopropyl]methylaniline trihydrochloride~~

Ammon.  
a<sup>8</sup>



Synthesis followed the procedure of Example 57. From 5-amino-2-(3,5-dimethoxy-4-methylthiophenyl)pyridine (182.0 mg, 0.660 mmol) synthesized by a similar procedure to that described in Reference Example 11, N,N-bis[2-[N-[2-(3,5-dimethoxy-4-methylthio-  
5 phenyl)-5-pyridyl]carbamoyl]ethyl]methylamine was obtained as a colorless viscous oil [136.0 mg, yield: 61% based on 5-amino-2-(3,5-dimethoxy-4-methylthiophenyl)pyridine]. From N,N-bis[2-[N-[2-(3,5-dimethoxy-4-methylthiophenyl)-5-pyridyl]carbamoyl]ethyl]methylamine (671.0 mg, 0.970 mmol), N,N-bis[N-[2-(3,5-dimethoxy-4-methylthiophenyl)-5-pyridyl]-3-aminopropyl]methylamine was obtained as a colorless oil (508.0 mg, yield: 79%). From N,N-bis[N-[2-(3,5-dimethoxy-4-methylthiophenyl)-5-pyridyl]-3-aminopropyl]methylamine (67.0 mg, 0.100 mmol), N,N-bis[N-[2-(3,5-dimethoxy-4-methylthiophenyl)-5-pyridyl]-N-methyl-3-aminopropyl]methylamine was obtained as a colorless oil (44.0 mg, yield: 64%). From N,N-bis[N-[2-(3,5-dimethoxy-4-methylthiophenyl)-5-pyridyl]-N-methyl-3-aminopropyl]methylamine (67.0 mg, 0.100 mmol), the title compound was obtained as a  
15 yellow amorphous powder (42.0 mg, yield: 52%).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>)(data for the free base of the title compound)

δ: 1.74-1.85(m, 4H), 2.23(s, 3H), 2.35-2.44(m, 4H),  
2.39(s, 6H), 3.01(s, 6H), 3.45(t, J=7.3Hz, 4H),  
20 3.98(s, 12H), 7.04(dd, J=2.9, 8.8Hz, 2H), 7.15(s, 4H), 7.59(d, J=8.8Hz, 2H), 8.22(d, J=2.9Hz, 2H).

Test 1 (Evaluation of IgE antibody production inhibiting activity)

From a mouse (Balb/C, male, 8 weeks old), the spleen was enucleated. The spleen was shredded in 10% FBS/RPMI 1640, and was then disintegrated into single cells through a 70-mesh screen. Those single cells were hemolyzed with Gey's solution, and using RPMI 1640 medium/25 mM HEPES/0.3% BSA, a spleen cell suspension ( $1 \times 10^7$  cells/mL) was prepared. After an aliquot of the suspension was reacted with a rat anti-mouse Thy-1.2 monoclonal antibody (product of Cedarlane Laboratories Limited) at 4°C for 1 hour, centrifugation was conducted. Precipitated cells were suspended again ( $1 \times 10^7$  cells/mL, RPMI/HEPES/BSA). After the suspension was next reacted with a low-cytotoxic rabbit complement (product of Cedarlane Laboratories Limited) at 37°C for 1 hour, dead cells were removed by specific gravity centrifugation using Lympholyte M (product of Cedarlane Laboratories Limited) so that a B cell fraction was obtained as viable cells.

Using a 96-well plate, the B cells ( $2 \times 10^5$  cells/0.2 mL/well) were incubated together with LPS (E. coli 026:B6, product of DIFCO Laboratories, Inc.) for 1 day. Mouse IL-4 (product of Genzyme Corp.) was then added, followed by further incubation for 6 days.

The IgE antibody production inhibiting activity of each drug was calculated by adding the drug Day 1 of the incubation and assaying the quantity of IgE in the culture supernatant by ELISA after the incubation. Inhibition activity ( $IC_{50}$ ) is presented in Table 1.

Further, the solubility (%) of each compound in water was also estimated. The results are presented in Table 1.

TABLE 1

Compound (Example No.)	IC <sub>50</sub> ( $\mu$ M)	Solubility (%)
7	0.04	5
14	0.04	0.1
30	0.10	0.1
36	0.03	10
37	0.10	10
39	0.08	10
41	0.05	0.1*
42	0.10	10
43	0.10	10*
44	0.10	10
47	0.05	0.1
49	0.04	10
57	0.10	1

\* Solubility in 1 M hydrochloric acid

#### Industrial Applicability

The bis(2-aryl-5-pyridyl) derivatives (1) of the present invention and salts thereof have excellent IgE antibody production inhibiting activity and are useful as medicinal agents for the prevention or treatment of allergic immune diseases.

Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the scope of the appended claims, the invention may be practiced otherwise than as specifically described herein.